

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, all ages				
...	All causes	2,813,503	100.0	863.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	647,457	23.0	198.8
2	Malignant neoplasms (C00-C97)	599,108	21.3	183.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169,936	6.0	52.2
4	Chronic lower respiratory diseases (J40-J47)	160,201	5.7	49.2
5	Cerebrovascular diseases (I60-I69)	146,383	5.2	44.9
6	Alzheimer disease (G30)	121,404	4.3	37.3
7	Diabetes mellitus (E10-E14)	83,564	3.0	25.7
8	Influenza and pneumonia (J09-J18)	55,672	2.0	17.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50,633	1.8	15.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47,173	1.7	14.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	41,743	1.5	12.8
12	Septicemia (A40-A41)	40,922	1.5	12.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35,316	1.3	10.8
14	Parkinson disease (G20-G21)	31,963	1.1	9.8
15	Pneumonitis due to solids and liquids (J69)	20,108	0.7	6.2
...	All other causes (residual)	561,920	20.0	172.5
All races and origins, both sexes, 1-4 years				
...	All causes	3,880	100.0	24.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,267	32.7	7.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	424	10.9	2.7
3	Malignant neoplasms (C00-C97)	325	8.4	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	303	7.8	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	3.3	0.8
6	Influenza and pneumonia (J09-J18)	104	2.7	0.7
7	Cerebrovascular diseases (I60-I69)	66	1.7	0.4
8	Septicemia (A40-A41)	48	1.2	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	1.1	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	42	1.1	0.3
11	Chronic lower respiratory diseases (J40-J47)	31	0.8	0.2
12	Anemias (D50-D64)	24	0.6	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	24	0.6	0.2
14	Acute bronchitis and bronchiolitis (J20-J21)	22	0.6	0.1
15	Meningitis (G00,G03)	19	0.5	*
...	All other causes (residual)	1,010	26.0	6.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 5-14 years				
...	All causes	5,571	100.0	13.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,578	28.3	3.8
2	Malignant neoplasms (C00-C97)	855	15.3	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	522	9.4	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	379	6.8	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	332	6.0	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	179	3.2	0.4
7	Chronic lower respiratory diseases (J40-J47)	134	2.4	0.3
8	Influenza and pneumonia (J09-J18)	113	2.0	0.3
9	Cerebrovascular diseases (I60-I69)	97	1.7	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	1.1	0.2
11	Septicemia (A40-A41)	57	1.0	0.1
12	Diabetes mellitus (E10-E14)	34	0.6	0.1
13	Complications of medical and surgical care (Y40-Y84,Y88)	25	0.4	0.1
14	Anemias (D50-D64)	24	0.4	0.1
15	Certain conditions originating in the perinatal period (P00-P96)	21	0.4	0.1
...	All other causes (residual)	1,159	20.8	2.8
All races and origins, both sexes, 15-24 years				
...	All causes	32,025	100.0	74.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,441	42.0	31.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,252	19.5	14.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,905	15.3	11.3
4	Malignant neoplasms (C00-C97)	1,374	4.3	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	913	2.9	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	355	1.1	0.8
7	Diabetes mellitus (E10-E14)	248	0.8	0.6
8	Influenza and pneumonia (J09-J18)	190	0.6	0.4
9	Chronic lower respiratory diseases (J40-J47)	188	0.6	0.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	168	0.5	0.4
11	Cerebrovascular diseases (I60-I69)	155	0.5	0.4
12	Septicemia (A40-A41)	123	0.4	0.3
13	Legal intervention (Y35,Y89.0)	97	0.3	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	69	0.2	0.2
15	Anemias (D50-D64)	69	0.2	0.2
...	All other causes (residual)	3,398	10.6	7.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 25-34 years				
...	All causes	60,215	100.0	132.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25,669	42.6	56.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,948	13.2	17.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,488	9.1	12.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,681	6.1	8.1
5	Malignant neoplasms (C00-C97)	3,616	6.0	8.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	918	1.5	2.0
7	Diabetes mellitus (E10-E14)	823	1.4	1.8
8	Cerebrovascular diseases (I60-I69)	593	1.0	1.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	513	0.9	1.1
10	Pregnancy, childbirth and the puerperium (O00-O99)	512	0.9	1.1
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	437	0.7	1.0
12	Influenza and pneumonia (J09-J18)	405	0.7	0.9
13	Septicemia (A40-A41)	392	0.7	0.9
14	Chronic lower respiratory diseases (J40-J47)	332	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	255	0.4	0.6
...	All other causes (residual)	8,633	14.3	19.0
All races and origins, both sexes, 35-44 years				
...	All causes	79,796	100.0	195.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,828	28.6	55.8
2	Malignant neoplasms (C00-C97)	10,900	13.7	26.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,401	13.0	25.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,335	9.2	17.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,351	4.2	8.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,000	3.8	7.3
7	Diabetes mellitus (E10-E14)	2,118	2.7	5.2
8	Cerebrovascular diseases (I60-I69)	1,811	2.3	4.4
9	Septicemia (A40-A41)	854	1.1	2.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	831	1.0	2.0
11	Influenza and pneumonia (J09-J18)	782	1.0	1.9
12	Chronic lower respiratory diseases (J40-J47)	713	0.9	1.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	692	0.9	1.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	514	0.6	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	419	0.5	1.0
...	All other causes (residual)	13,247	16.6	32.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 45-54 years				
...	All causes	170,142	100.0	401.5
1	Malignant neoplasms (C00-C97)	39,266	23.1	92.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,658	19.2	77.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24,461	14.4	57.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,561	5.0	20.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	8,312	4.9	19.6
6	Diabetes mellitus (E10-E14)	6,409	3.8	15.1
7	Cerebrovascular diseases (I60-I69)	5,198	3.1	12.3
8	Chronic lower respiratory diseases (J40-J47)	3,975	2.3	9.4
9	Septicemia (A40-A41)	2,441	1.4	5.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,275	1.3	5.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,197	1.3	5.2
12	Influenza and pneumonia (J09-J18)	2,013	1.2	4.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,642	1.0	3.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,609	0.9	3.8
15	Viral hepatitis (B15-B19)	927	0.5	2.2
...	All other causes (residual)	28,198	16.6	66.5
All races and origins, both sexes, 55-64 years				
...	All causes	372,006	100.0	885.8
1	Malignant neoplasms (C00-C97)	114,810	30.9	273.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	80,102	21.5	190.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23,408	6.3	55.7
4	Chronic lower respiratory diseases (J40-J47)	18,667	5.0	44.4
5	Diabetes mellitus (E10-E14)	14,904	4.0	35.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	13,737	3.7	32.7
7	Cerebrovascular diseases (I60-I69)	12,708	3.4	30.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,982	2.1	19.0
9	Septicemia (A40-A41)	5,838	1.6	13.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,671	1.5	13.5
11	Influenza and pneumonia (J09-J18)	5,045	1.4	12.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,991	1.1	9.5
13	Viral hepatitis (B15-B19)	2,517	0.7	6.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,737	0.5	4.1
15	Pneumonitis due to solids and liquids (J69)	1,610	0.4	3.8
...	All other causes (residual)	59,279	15.9	141.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 65-74 years				
...	All causes	531,610	100.0	1,790.9
1	Malignant neoplasms (C00-C97)	168,458	31.7	567.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	116,633	21.9	392.9
3	Chronic lower respiratory diseases (J40-J47)	39,704	7.5	133.8
4	Cerebrovascular diseases (I60-I69)	22,690	4.3	76.4
5	Diabetes mellitus (E10-E14)	21,344	4.0	71.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,038	2.8	50.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,302	1.9	34.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	9,400	1.8	31.7
9	Septicemia (A40-A41)	9,164	1.7	30.9
10	Influenza and pneumonia (J09-J18)	8,784	1.7	29.6
11	Alzheimer disease (G30)	7,281	1.4	24.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,856	1.1	19.7
13	Parkinson disease (G20-G21)	4,661	0.9	15.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,620	0.9	15.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,108	0.6	10.5
...	All other causes (residual)	84,567	15.9	284.9
All races and origins, both sexes, 75-84 years				
...	All causes	657,759	100.0	4,472.6
1	Malignant neoplasms (C00-C97)	155,917	23.7	1,060.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	151,249	23.0	1,028.4
3	Chronic lower respiratory diseases (J40-J47)	51,116	7.8	347.6
4	Cerebrovascular diseases (I60-I69)	38,697	5.9	263.1
5	Alzheimer disease (G30)	32,309	4.9	219.7
6	Diabetes mellitus (E10-E14)	20,703	3.1	140.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,661	2.5	113.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,093	2.1	95.8
9	Influenza and pneumonia (J09-J18)	13,802	2.1	93.8
10	Parkinson disease (G20-G21)	13,172	2.0	89.6
11	Septicemia (A40-A41)	10,648	1.6	72.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,211	1.2	55.8
13	Pneumonitis due to solids and liquids (J69)	5,236	0.8	35.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,735	0.7	32.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,610	0.7	31.3
...	All other causes (residual)	116,600	17.7	792.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 85 years and over				
...	All causes	878,035	100.0	13,573.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	251,170	28.6	3,882.9
2	Malignant neoplasms (C00-C97)	103,521	11.8	1,600.3
3	Alzheimer disease (G30)	80,517	9.2	1,244.7
4	Cerebrovascular diseases (I60-I69)	64,266	7.3	993.5
5	Chronic lower respiratory diseases (J40-J47)	45,319	5.2	700.6
6	Influenza and pneumonia (J09-J18)	24,276	2.8	375.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24,252	2.8	374.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,275	2.0	267.1
9	Diabetes mellitus (E10-E14)	16,973	1.9	262.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14,969	1.7	231.4
11	Parkinson disease (G20-G21)	13,344	1.5	206.3
12	Septicemia (A40-A41)	11,210	1.3	173.3
13	Pneumonitis due to solids and liquids (J69)	9,361	1.1	144.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,312	0.6	82.1
15	Nutritional deficiencies (E40-E64)	4,294	0.5	66.4
...	All other causes (residual)	191,976	21.9	2,967.8
All races and origins, male, all ages				
...	All causes	1,439,111	100.0	897.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	347,879	24.2	216.9
2	Malignant neoplasms (C00-C97)	315,147	21.9	196.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109,722	7.6	68.4
4	Chronic lower respiratory diseases (J40-J47)	75,005	5.2	46.8
5	Cerebrovascular diseases (I60-I69)	61,645	4.3	38.4
6	Diabetes mellitus (E10-E14)	46,302	3.2	28.9
7	Alzheimer disease (G30)	37,325	2.6	23.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36,782	2.6	22.9
9	Influenza and pneumonia (J09-J18)	26,558	1.8	16.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	26,451	1.8	16.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25,744	1.8	16.0
12	Septicemia (A40-A41)	19,603	1.4	12.2
13	Parkinson disease (G20-G21)	19,388	1.3	12.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15,749	1.1	9.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15,524	1.1	9.7
...	All other causes (residual)	260,287	18.1	162.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 1-4 years				
...	All causes	2,232	100.0	27.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	770	34.5	9.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	225	10.1	2.8
3	Malignant neoplasms (C00-C97)	184	8.2	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	7.5	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	3.0	0.8
6	Influenza and pneumonia (J09-J18)	63	2.8	0.8
7	Cerebrovascular diseases (I60-I69)	42	1.9	0.5
8	Septicemia (A40-A41)	26	1.2	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.9	*
9	Certain conditions originating in the perinatal period (P00-P96)	19	0.9	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	16	0.7	*
13	Acute bronchitis and bronchiolitis (J20-J21)	15	0.7	*
14	Meningitis (G00,G03)	14	0.6	*
15	Anemias (D50-D64)	8	0.4	*
...	All other causes (residual)	579	25.9	7.1
All races and origins, male, 5-14 years				
...	All causes	3,269	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	985	30.1	4.7
2	Malignant neoplasms (C00-C97)	455	13.9	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	352	10.8	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	208	6.4	1.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	196	6.0	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	2.9	0.5
7	Chronic lower respiratory diseases (J40-J47)	84	2.6	0.4
8	Cerebrovascular diseases (I60-I69)	53	1.6	0.3
9	Influenza and pneumonia (J09-J18)	49	1.5	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	1.1	0.2
11	Septicemia (A40-A41)	31	0.9	0.1
12	Diabetes mellitus (E10-E14)	20	0.6	0.1
13	Anemias (D50-D64)	15	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	12	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.4	*
...	All other causes (residual)	666	20.4	3.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 15-24 years				
...	All causes	23,503	100.0	106.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,746	41.5	44.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,027	21.4	22.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,234	18.0	19.1
4	Malignant neoplasms (C00-C97)	833	3.5	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	620	2.6	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	208	0.9	0.9
7	Diabetes mellitus (E10-E14)	144	0.6	0.7
8	Chronic lower respiratory diseases (J40-J47)	115	0.5	0.5
9	Influenza and pneumonia (J09-J18)	109	0.5	0.5
10	Legal intervention (Y35,Y89.0)	90	0.4	0.4
11	Cerebrovascular diseases (I60-I69)	83	0.4	0.4
12	Septicemia (A40-A41)	62	0.3	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	52	0.2	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	30	0.1	0.1
...	All other causes (residual)	2,105	9.0	9.5
All races and origins, male, 25-34 years				
...	All causes	42,149	100.0	183.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,001	45.1	82.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,348	15.1	27.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,607	10.9	20.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,449	5.8	10.7
5	Malignant neoplasms (C00-C97)	1,690	4.0	7.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	551	1.3	2.4
7	Diabetes mellitus (E10-E14)	471	1.1	2.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	369	0.9	1.6
9	Cerebrovascular diseases (I60-I69)	339	0.8	1.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	276	0.7	1.2
11	Influenza and pneumonia (J09-J18)	215	0.5	0.9
12	Septicemia (A40-A41)	200	0.5	0.9
13	Chronic lower respiratory diseases (J40-J47)	192	0.5	0.8
14	Legal intervention (Y35,Y89.0)	181	0.4	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	142	0.3	0.6
...	All other causes (residual)	5,118	12.1	22.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 35-44 years				
...	All causes	50,792	100.0	249.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,243	32.0	79.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,074	13.9	34.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,596	11.0	27.5
4	Malignant neoplasms (C00-C97)	4,725	9.3	23.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,676	5.3	13.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,853	3.6	9.1
7	Diabetes mellitus (E10-E14)	1,298	2.6	6.4
8	Cerebrovascular diseases (I60-I69)	1,034	2.0	5.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	567	1.1	2.8
10	Septicemia (A40-A41)	456	0.9	2.2
11	Influenza and pneumonia (J09-J18)	437	0.9	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	393	0.8	1.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	354	0.7	1.7
14	Chronic lower respiratory diseases (J40-J47)	323	0.6	1.6
15	Aortic aneurysm and dissection (I71)	249	0.5	1.2
...	All other causes (residual)	7,514	14.8	36.9
All races and origins, male, 45-54 years				
...	All causes	103,804	100.0	496.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,808	22.0	109.1
2	Malignant neoplasms (C00-C97)	18,898	18.2	90.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,877	16.3	80.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,331	6.1	30.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,389	5.2	25.8
6	Diabetes mellitus (E10-E14)	4,020	3.9	19.2
7	Cerebrovascular diseases (I60-I69)	3,010	2.9	14.4
8	Chronic lower respiratory diseases (J40-J47)	1,832	1.8	8.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,737	1.7	8.3
10	Septicemia (A40-A41)	1,323	1.3	6.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,288	1.2	6.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,167	1.1	5.6
13	Influenza and pneumonia (J09-J18)	1,076	1.0	5.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,061	1.0	5.1
15	Viral hepatitis (B15-B19)	624	0.6	3.0
...	All other causes (residual)	16,363	15.8	78.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 55-64 years				
...	All causes	225,335	100.0	1,112.3
1	Malignant neoplasms (C00-C97)	62,380	27.7	307.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,335	24.6	273.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,260	7.2	80.3
4	Chronic lower respiratory diseases (J40-J47)	9,424	4.2	46.5
5	Diabetes mellitus (E10-E14)	9,282	4.1	45.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	9,223	4.1	45.5
7	Cerebrovascular diseases (I60-I69)	7,332	3.3	36.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,040	2.7	29.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,199	1.4	15.8
10	Septicemia (A40-A41)	3,160	1.4	15.6
11	Influenza and pneumonia (J09-J18)	2,826	1.3	14.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,473	1.1	12.2
13	Viral hepatitis (B15-B19)	1,739	0.8	8.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,314	0.6	6.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,143	0.5	5.6
...	All other causes (residual)	34,205	15.2	168.8
All races and origins, male, 65-74 years				
...	All causes	303,931	100.0	2,190.2
1	Malignant neoplasms (C00-C97)	93,082	30.6	670.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,723	24.6	538.5
3	Chronic lower respiratory diseases (J40-J47)	20,183	6.6	145.4
4	Diabetes mellitus (E10-E14)	12,602	4.1	90.8
5	Cerebrovascular diseases (I60-I69)	12,032	4.0	86.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,797	3.2	70.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,059	2.0	43.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,642	1.9	40.7
9	Influenza and pneumonia (J09-J18)	4,782	1.6	34.5
10	Septicemia (A40-A41)	4,635	1.5	33.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,638	1.2	26.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,236	1.1	23.3
13	Alzheimer disease (G30)	3,085	1.0	22.2
14	Parkinson disease (G20-G21)	3,072	1.0	22.1
15	Pneumonitis due to solids and liquids (J69)	1,876	0.6	13.5
...	All other causes (residual)	45,487	15.0	327.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 75-84 years				
...	All causes	336,671	100.0	5,254.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	83,738	24.9	1,306.8
2	Malignant neoplasms (C00-C97)	83,349	24.8	1,300.7
3	Chronic lower respiratory diseases (J40-J47)	24,139	7.2	376.7
4	Cerebrovascular diseases (I60-I69)	17,524	5.2	273.5
5	Alzheimer disease (G30)	12,098	3.6	188.8
6	Diabetes mellitus (E10-E14)	11,211	3.3	175.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,267	2.8	144.6
8	Parkinson disease (G20-G21)	8,461	2.5	132.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,223	2.1	112.7
10	Influenza and pneumonia (J09-J18)	7,056	2.1	110.1
11	Septicemia (A40-A41)	5,068	1.5	79.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,774	1.1	58.9
13	Pneumonitis due to solids and liquids (J69)	3,108	0.9	48.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,619	0.8	40.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,546	0.8	39.7
...	All other causes (residual)	55,490	16.5	866.0
All races and origins, male, 85 years and over				
...	All causes	334,866	100.0	14,689.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	100,787	30.1	4,421.1
2	Malignant neoplasms (C00-C97)	49,519	14.8	2,172.2
3	Alzheimer disease (G30)	21,574	6.4	946.4
4	Cerebrovascular diseases (I60-I69)	20,137	6.0	883.3
5	Chronic lower respiratory diseases (J40-J47)	18,682	5.6	819.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,001	3.0	438.7
7	Influenza and pneumonia (J09-J18)	9,846	2.9	431.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,779	2.3	341.2
9	Parkinson disease (G20-G21)	7,339	2.2	321.9
10	Diabetes mellitus (E10-E14)	7,251	2.2	318.1
11	Pneumonitis due to solids and liquids (J69)	4,833	1.4	212.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,752	1.4	208.5
13	Septicemia (A40-A41)	4,563	1.4	200.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,511	0.7	110.1
15	Nutritional deficiencies (E40-E64)	1,301	0.4	57.1
...	All other causes (residual)	63,991	19.1	2,807.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, all ages				
...	All causes	1,374,392	100.0	831.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	299,578	21.8	181.2
2	Malignant neoplasms (C00-C97)	283,961	20.7	171.8
3	Chronic lower respiratory diseases (J40-J47)	85,196	6.2	51.5
4	Cerebrovascular diseases (I60-I69)	84,738	6.2	51.3
5	Alzheimer disease (G30)	84,079	6.1	50.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60,214	4.4	36.4
7	Diabetes mellitus (E10-E14)	37,262	2.7	22.5
8	Influenza and pneumonia (J09-J18)	29,114	2.1	17.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24,889	1.8	15.1
10	Septicemia (A40-A41)	21,319	1.6	12.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19,567	1.4	11.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	15,292	1.1	9.3
13	Parkinson disease (G20-G21)	12,575	0.9	7.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,391	0.8	6.3
15	Pneumonitis due to solids and liquids (J69)	8,758	0.6	5.3
...	All other causes (residual)	297,459	21.6	179.9
All races and origins, female, 1-4 years				
...	All causes	1,648	100.0	21.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	497	30.2	6.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	199	12.1	2.5
3	Malignant neoplasms (C00-C97)	141	8.6	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	135	8.2	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	3.6	0.8
6	Influenza and pneumonia (J09-J18)	41	2.5	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.5	0.3
8	Cerebrovascular diseases (I60-I69)	24	1.5	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	23	1.4	0.3
10	Septicemia (A40-A41)	22	1.3	0.3
11	Anemias (D50-D64)	16	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	15	0.9	*
13	Acute bronchitis and bronchiolitis (J20-J21)	7	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
15	Meningitis (G00,G03)	5	0.3	*
...	All other causes (residual)	431	26.2	5.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 5-14 years				
...	All causes	2,302	100.0	11.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	593	25.8	2.9
2	Malignant neoplasms (C00-C97)	400	17.4	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	171	7.4	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	170	7.4	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	136	5.9	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	3.6	0.4
7	Influenza and pneumonia (J09-J18)	64	2.8	0.3
8	Chronic lower respiratory diseases (J40-J47)	50	2.2	0.2
9	Cerebrovascular diseases (I60-I69)	44	1.9	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	1.2	0.1
11	Septicemia (A40-A41)	26	1.1	0.1
12	Diabetes mellitus (E10-E14)	14	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.6	*
14	Anemias (D50-D64)	9	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	9	0.4	*
...	All other causes (residual)	493	21.4	2.5
All races and origins, female, 15-24 years				
...	All causes	8,522	100.0	40.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,695	43.4	17.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,225	14.4	5.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	671	7.9	3.2
4	Malignant neoplasms (C00-C97)	541	6.3	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	293	3.4	1.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	168	2.0	0.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	147	1.7	0.7
8	Diabetes mellitus (E10-E14)	104	1.2	0.5
9	Influenza and pneumonia (J09-J18)	81	1.0	0.4
10	Chronic lower respiratory diseases (J40-J47)	73	0.9	0.3
11	Cerebrovascular diseases (I60-I69)	72	0.8	0.3
12	Septicemia (A40-A41)	61	0.7	0.3
13	Anemias (D50-D64)	40	0.5	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.4	0.2
15	Complications of medical and surgical care (Y40-Y84,Y88)	26	0.3	0.1
...	All other causes (residual)	1,290	15.1	6.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 25-34 years				
...	All causes	18,066	100.0	80.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,668	36.9	29.8
2	Malignant neoplasms (C00-C97)	1,926	10.7	8.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,600	8.9	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,232	6.8	5.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	881	4.9	3.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	512	2.8	2.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	367	2.0	1.6
8	Diabetes mellitus (E10-E14)	352	1.9	1.6
9	Cerebrovascular diseases (I60-I69)	254	1.4	1.1
10	Septicemia (A40-A41)	192	1.1	0.9
11	Influenza and pneumonia (J09-J18)	190	1.1	0.9
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	161	0.9	0.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	144	0.8	0.6
14	Chronic lower respiratory diseases (J40-J47)	140	0.8	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	113	0.6	0.5
...	All other causes (residual)	3,334	18.5	14.9
All races and origins, female, 35-44 years				
...	All causes	29,004	100.0	141.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,585	22.7	32.1
2	Malignant neoplasms (C00-C97)	6,175	21.3	30.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,327	11.5	16.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,739	6.0	8.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,147	4.0	5.6
6	Diabetes mellitus (E10-E14)	820	2.8	4.0
7	Cerebrovascular diseases (I60-I69)	777	2.7	3.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	675	2.3	3.3
9	Septicemia (A40-A41)	398	1.4	1.9
10	Chronic lower respiratory diseases (J40-J47)	390	1.3	1.9
11	Influenza and pneumonia (J09-J18)	345	1.2	1.7
12	Pregnancy, childbirth and the puerperium (O00-O99)	328	1.1	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	299	1.0	1.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	264	0.9	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	190	0.7	0.9
...	All other causes (residual)	5,545	19.1	27.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 45-54 years				
...	All causes	66,338	100.0	309.0
1	Malignant neoplasms (C00-C97)	20,368	30.7	94.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,850	14.8	45.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,584	11.4	35.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,923	4.4	13.6
5	Diabetes mellitus (E10-E14)	2,389	3.6	11.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,230	3.4	10.4
7	Cerebrovascular diseases (I60-I69)	2,188	3.3	10.2
8	Chronic lower respiratory diseases (J40-J47)	2,143	3.2	10.0
9	Septicemia (A40-A41)	1,118	1.7	5.2
10	Influenza and pneumonia (J09-J18)	937	1.4	4.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	909	1.4	4.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	581	0.9	2.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	538	0.8	2.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	442	0.7	2.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	352	0.5	1.6
...	All other causes (residual)	11,786	17.8	54.9
All races and origins, female, 55-64 years				
...	All causes	146,671	100.0	674.7
1	Malignant neoplasms (C00-C97)	52,430	35.7	241.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,767	16.9	113.9
3	Chronic lower respiratory diseases (J40-J47)	9,243	6.3	42.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,148	4.9	32.9
5	Diabetes mellitus (E10-E14)	5,622	3.8	25.9
6	Cerebrovascular diseases (I60-I69)	5,376	3.7	24.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,514	3.1	20.8
8	Septicemia (A40-A41)	2,678	1.8	12.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,472	1.7	11.4
10	Influenza and pneumonia (J09-J18)	2,219	1.5	10.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,942	1.3	8.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,518	1.0	7.0
13	Viral hepatitis (B15-B19)	778	0.5	3.6
14	Alzheimer disease (G30)	665	0.5	3.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	633	0.4	2.9
...	All other causes (residual)	24,666	16.8	113.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 65-74 years				
...	All causes	227,679	100.0	1,440.4
1	Malignant neoplasms (C00-C97)	75,376	33.1	476.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,910	18.4	265.1
3	Chronic lower respiratory diseases (J40-J47)	19,521	8.6	123.5
4	Cerebrovascular diseases (I60-I69)	10,658	4.7	67.4
5	Diabetes mellitus (E10-E14)	8,742	3.8	55.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,241	2.3	33.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,660	2.0	29.5
8	Septicemia (A40-A41)	4,529	2.0	28.7
9	Alzheimer disease (G30)	4,196	1.8	26.5
10	Influenza and pneumonia (J09-J18)	4,002	1.8	25.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	3,341	1.5	21.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,620	1.2	16.6
13	Parkinson disease (G20-G21)	1,589	0.7	10.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,251	0.5	7.9
15	Pneumonitis due to solids and liquids (J69)	1,116	0.5	7.1
...	All other causes (residual)	38,927	17.1	246.3
All races and origins, female, 75-84 years				
...	All causes	321,088	100.0	3,869.1
1	Malignant neoplasms (C00-C97)	72,568	22.6	874.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	67,511	21.0	813.5
3	Chronic lower respiratory diseases (J40-J47)	26,977	8.4	325.1
4	Cerebrovascular diseases (I60-I69)	21,173	6.6	255.1
5	Alzheimer disease (G30)	20,211	6.3	243.5
6	Diabetes mellitus (E10-E14)	9,492	3.0	114.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,394	2.3	89.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,870	2.1	82.8
9	Influenza and pneumonia (J09-J18)	6,746	2.1	81.3
10	Septicemia (A40-A41)	5,580	1.7	67.2
11	Parkinson disease (G20-G21)	4,711	1.5	56.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,437	1.4	53.5
13	Pneumonitis due to solids and liquids (J69)	2,128	0.7	25.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,116	0.7	25.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,064	0.6	24.9
...	All other causes (residual)	61,110	19.0	736.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 85 years and over				
...	All causes	543,169	100.0	12,966.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	150,383	27.7	3,589.9
2	Alzheimer disease (G30)	58,943	10.9	1,407.1
3	Malignant neoplasms (C00-C97)	54,002	9.9	1,289.1
4	Cerebrovascular diseases (I60-I69)	44,129	8.1	1,053.4
5	Chronic lower respiratory diseases (J40-J47)	26,637	4.9	635.9
6	Influenza and pneumonia (J09-J18)	14,430	2.7	344.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,251	2.6	340.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,217	1.9	243.9
9	Diabetes mellitus (E10-E14)	9,722	1.8	232.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,496	1.7	226.7
11	Septicemia (A40-A41)	6,647	1.2	158.7
12	Parkinson disease (G20-G21)	6,005	1.1	143.4
13	Pneumonitis due to solids and liquids (J69)	4,528	0.8	108.1
14	Nutritional deficiencies (E40-E64)	2,993	0.6	71.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,801	0.5	66.9
...	All other causes (residual)	127,985	23.6	3,055.3
Hispanic, both sexes, all ages				
...	All causes	197,249	100.0	334.6
1	Malignant neoplasms (C00-C97)	40,550	20.6	68.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,402	20.0	66.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,757	8.5	28.4
4	Cerebrovascular diseases (I60-I69)	10,775	5.5	18.3
5	Diabetes mellitus (E10-E14)	9,216	4.7	15.6
6	Alzheimer disease (G30)	7,383	3.7	12.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,286	3.2	10.7
8	Chronic lower respiratory diseases (J40-J47)	5,599	2.8	9.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,933	2.0	6.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,928	2.0	6.7
11	Influenza and pneumonia (J09-J18)	3,851	2.0	6.5
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,186	1.6	5.4
13	Septicemia (A40-A41)	2,869	1.5	4.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,799	1.4	4.7
15	Certain conditions originating in the perinatal period (P00-P96)	2,409	1.2	4.1
...	All other causes (residual)	38,306	19.4	65.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 1-4 years				
...	All causes	805	100.0	19.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	29.1	5.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	13.9	2.7
3	Malignant neoplasms (C00-C97)	92	11.4	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	6.2	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	3.2	0.6
6	Influenza and pneumonia (J09-J18)	24	3.0	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.7	*
8	Cerebrovascular diseases (I60-I69)	12	1.5	*
9	Septicemia (A40-A41)	11	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	11	1.4	*
11	Acute bronchitis and bronchiolitis (J20-J21)	6	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.6	*
14	Meningitis (G00,G03)	4	0.5	*
15	Anemias (D50-D64)	3	0.4	*
...	All other causes (residual)	196	24.3	4.7
Hispanic, both sexes, 5-14 years				
...	All causes	1,165	100.0	11.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	302	25.9	2.9
2	Malignant neoplasms (C00-C97)	222	19.1	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	7.8	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	7.0	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	5.2	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	3.3	0.4
7	Cerebrovascular diseases (I60-I69)	24	2.1	0.2
8	Influenza and pneumonia (J09-J18)	21	1.8	0.2
9	Chronic lower respiratory diseases (J40-J47)	19	1.6	*
10	Septicemia (A40-A41)	18	1.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.2	*
12	Diabetes mellitus (E10-E14)	5	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
15	Diseases of appendix (K35-K38)	4	0.3	*
...	All other causes (residual)	256	22.0	2.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-24 years				
...	All causes	5,854	100.0	60.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,486	42.5	25.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	980	16.7	10.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	971	16.6	10.0
4	Malignant neoplasms (C00-C97)	371	6.3	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	131	2.2	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	1.2	0.7
7	Cerebrovascular diseases (I60-I69)	41	0.7	0.4
8	Pregnancy, childbirth and the puerperium (O00-O99)	34	0.6	0.3
9	Influenza and pneumonia (J09-J18)	32	0.5	0.3
10	Chronic lower respiratory diseases (J40-J47)	30	0.5	0.3
11	Diabetes mellitus (E10-E14)	29	0.5	0.3
12	Legal intervention (Y35,Y89.0)	27	0.5	0.3
13	Septicemia (A40-A41)	23	0.4	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.2	*
...	All other causes (residual)	601	10.3	6.2
Hispanic, both sexes, 25-34 years				
...	All causes	8,492	100.0	91.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,379	39.8	36.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	964	11.4	10.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	909	10.7	9.8
4	Malignant neoplasms (C00-C97)	728	8.6	7.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	495	5.8	5.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	173	2.0	1.9
7	Cerebrovascular diseases (I60-I69)	125	1.5	1.3
8	Diabetes mellitus (E10-E14)	113	1.3	1.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	84	1.0	0.9
10	Pregnancy, childbirth and the puerperium (O00-O99)	80	0.9	0.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	0.8	0.7
12	Septicemia (A40-A41)	59	0.7	0.6
13	Influenza and pneumonia (J09-J18)	55	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	0.6	0.6
15	Chronic lower respiratory diseases (J40-J47)	48	0.6	0.5
...	All other causes (residual)	1,161	13.7	12.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-44 years				
...	All causes	10,738	100.0	126.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,904	27.0	34.3
2	Malignant neoplasms (C00-C97)	1,777	16.5	21.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,165	10.8	13.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	704	6.6	8.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	627	5.8	7.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	580	5.4	6.8
7	Cerebrovascular diseases (I60-I69)	351	3.3	4.1
8	Diabetes mellitus (E10-E14)	280	2.6	3.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	145	1.4	1.7
10	Septicemia (A40-A41)	124	1.2	1.5
11	Influenza and pneumonia (J09-J18)	107	1.0	1.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	82	0.8	1.0
13	Chronic lower respiratory diseases (J40-J47)	67	0.6	0.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	57	0.5	0.7
15	Pregnancy, childbirth and the puerperium (O00-O99)	56	0.5	0.7
...	All other causes (residual)	1,712	15.9	20.2
Hispanic, both sexes, 45-54 years				
...	All causes	18,409	100.0	267.0
1	Malignant neoplasms (C00-C97)	4,409	24.0	63.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,951	16.0	42.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,643	14.4	38.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,465	8.0	21.2
5	Diabetes mellitus (E10-E14)	934	5.1	13.5
6	Cerebrovascular diseases (I60-I69)	733	4.0	10.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	582	3.2	8.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	318	1.7	4.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	293	1.6	4.2
10	Septicemia (A40-A41)	275	1.5	4.0
11	Influenza and pneumonia (J09-J18)	253	1.4	3.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	223	1.2	3.2
13	Viral hepatitis (B15-B19)	181	1.0	2.6
14	Chronic lower respiratory diseases (J40-J47)	168	0.9	2.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	167	0.9	2.4
...	All other causes (residual)	2,814	15.3	40.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-64 years				
...	All causes	28,533	100.0	606.8
1	Malignant neoplasms (C00-C97)	8,253	28.9	175.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,583	19.6	118.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,897	6.6	40.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,784	6.3	37.9
5	Diabetes mellitus (E10-E14)	1,742	6.1	37.0
6	Cerebrovascular diseases (I60-I69)	1,311	4.6	27.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	598	2.1	12.7
8	Chronic lower respiratory diseases (J40-J47)	546	1.9	11.6
9	Septicemia (A40-A41)	465	1.6	9.9
10	Influenza and pneumonia (J09-J18)	435	1.5	9.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	395	1.4	8.4
12	Viral hepatitis (B15-B19)	374	1.3	8.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	328	1.1	7.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	220	0.8	4.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	172	0.6	3.7
...	All other causes (residual)	4,430	15.5	94.2
Hispanic, both sexes, 65-74 years				
...	All causes	33,368	100.0	1,308.7
1	Malignant neoplasms (C00-C97)	9,951	29.8	390.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,147	21.4	280.3
3	Diabetes mellitus (E10-E14)	2,263	6.8	88.8
4	Cerebrovascular diseases (I60-I69)	1,714	5.1	67.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,249	3.7	49.0
6	Chronic lower respiratory diseases (J40-J47)	1,034	3.1	40.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	989	3.0	38.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	899	2.7	35.3
9	Influenza and pneumonia (J09-J18)	635	1.9	24.9
10	Septicemia (A40-A41)	583	1.7	22.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	510	1.5	20.0
12	Alzheimer disease (G30)	480	1.4	18.8
13	Parkinson disease (G20-G21)	270	0.8	10.6
14	Viral hepatitis (B15-B19)	175	0.5	6.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	160	0.5	6.3
...	All other causes (residual)	5,309	15.9	208.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-84 years				
...	All causes	39,830	100.0	3,379.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,379	23.5	795.8
2	Malignant neoplasms (C00-C97)	9,096	22.8	771.8
3	Cerebrovascular diseases (I60-I69)	2,757	6.9	233.9
4	Diabetes mellitus (E10-E14)	2,312	5.8	196.2
5	Alzheimer disease (G30)	2,008	5.0	170.4
6	Chronic lower respiratory diseases (J40-J47)	1,686	4.2	143.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,008	2.5	85.5
8	Influenza and pneumonia (J09-J18)	959	2.4	81.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	911	2.3	77.3
10	Parkinson disease (G20-G21)	755	1.9	64.1
11	Septicemia (A40-A41)	664	1.7	56.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	661	1.7	56.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	638	1.6	54.1
14	Pneumonitis due to solids and liquids (J69)	295	0.7	25.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	232	0.6	19.7
...	All other causes (residual)	6,469	16.2	548.9
Hispanic, both sexes, 85 years and over				
...	All causes	45,236	100.0	9,504.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,430	27.5	2,611.6
2	Malignant neoplasms (C00-C97)	5,634	12.5	1,183.7
3	Alzheimer disease (G30)	4,778	10.6	1,003.9
4	Cerebrovascular diseases (I60-I69)	3,695	8.2	776.3
5	Chronic lower respiratory diseases (J40-J47)	1,991	4.4	418.3
6	Diabetes mellitus (E10-E14)	1,537	3.4	322.9
7	Influenza and pneumonia (J09-J18)	1,295	2.9	272.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	991	2.2	208.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	965	2.1	202.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	918	2.0	192.9
11	Parkinson disease (G20-G21)	725	1.6	152.3
12	Septicemia (A40-A41)	616	1.4	129.4
13	Pneumonitis due to solids and liquids (J69)	388	0.9	81.5
14	Nutritional deficiencies (E40-E64)	254	0.6	53.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	229	0.5	48.1
...	All other causes (residual)	8,790	19.4	1,846.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	108,579	100.0	364.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,009	20.3	73.9
2	Malignant neoplasms (C00-C97)	21,034	19.4	70.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,455	11.5	41.8
4	Cerebrovascular diseases (I60-I69)	5,073	4.7	17.0
5	Diabetes mellitus (E10-E14)	5,054	4.7	17.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,369	4.0	14.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,175	2.9	10.7
8	Chronic lower respiratory diseases (J40-J47)	2,759	2.5	9.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,588	2.4	8.7
10	Alzheimer disease (G30)	2,297	2.1	7.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,988	1.8	6.7
12	Influenza and pneumonia (J09-J18)	1,911	1.8	6.4
13	Septicemia (A40-A41)	1,473	1.4	4.9
14	Certain conditions originating in the perinatal period (P00-P96)	1,356	1.2	4.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,354	1.2	4.5
...	All other causes (residual)	19,684	18.1	66.1
Hispanic, male, 1-4 years				
...	All causes	455	100.0	21.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	31.4	6.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	12.1	2.6
3	Malignant neoplasms (C00-C97)	50	11.0	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	6.6	1.4
5	Influenza and pneumonia (J09-J18)	15	3.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	3.1	*
7	Cerebrovascular diseases (I60-I69)	9	2.0	*
8	Septicemia (A40-A41)	7	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.3	*
11	Acute bronchitis and bronchiolitis (J20-J21)	5	1.1	*
12	Chronic lower respiratory diseases (J40-J47)	4	0.9	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.9	*
14	Meningitis (G00,G03)	2	0.4	*
14	Diseases of appendix (K35-K38)	2	0.4	*
...	All other causes (residual)	102	22.4	4.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-14 years				
...	All causes	677	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	26.7	3.4
2	Malignant neoplasms (C00-C97)	113	16.7	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	9.0	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	6.9	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	5.6	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	3.8	0.5
7	Cerebrovascular diseases (I60-I69)	10	1.5	*
7	Influenza and pneumonia (J09-J18)	10	1.5	*
7	Chronic lower respiratory diseases (J40-J47)	10	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.3	*
11	Septicemia (A40-A41)	8	1.2	*
12	Diabetes mellitus (E10-E14)	5	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
15	Diseases of appendix (K35-K38)	2	0.3	*
...	All other causes (residual)	150	22.2	2.8
Hispanic, male, 15-24 years				
...	All causes	4,387	100.0	87.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,890	43.1	37.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	839	19.1	16.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	780	17.8	15.6
4	Malignant neoplasms (C00-C97)	232	5.3	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	99	2.3	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.9	0.8
7	Cerebrovascular diseases (I60-I69)	26	0.6	0.5
8	Legal intervention (Y35,Y89.0)	25	0.6	0.5
9	Diabetes mellitus (E10-E14)	19	0.4	*
10	Influenza and pneumonia (J09-J18)	16	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	16	0.4	*
12	Septicemia (A40-A41)	15	0.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.1	*
15	Anemias (D50-D64)	6	0.1	*
...	All other causes (residual)	362	8.3	7.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-34 years				
...	All causes	6,309	100.0	129.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,727	43.2	55.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	796	12.6	16.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	759	12.0	15.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	382	6.1	7.8
5	Malignant neoplasms (C00-C97)	351	5.6	7.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	121	1.9	2.5
7	Cerebrovascular diseases (I60-I69)	75	1.2	1.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	70	1.1	1.4
9	Diabetes mellitus (E10-E14)	69	1.1	1.4
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	0.7	0.9
11	Legal intervention (Y35,Y89.0)	40	0.6	0.8
12	Septicemia (A40-A41)	35	0.6	0.7
12	Influenza and pneumonia (J09-J18)	35	0.6	0.7
14	Chronic lower respiratory diseases (J40-J47)	32	0.5	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.5	0.6
...	All other causes (residual)	740	11.7	15.1
Hispanic, male, 35-44 years				
...	All causes	7,358	100.0	169.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,335	31.7	53.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	856	11.6	19.7
3	Malignant neoplasms (C00-C97)	748	10.2	17.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	569	7.7	13.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	465	6.3	10.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	463	6.3	10.6
7	Cerebrovascular diseases (I60-I69)	241	3.3	5.5
8	Diabetes mellitus (E10-E14)	187	2.5	4.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	117	1.6	2.7
10	Septicemia (A40-A41)	75	1.0	1.7
11	Influenza and pneumonia (J09-J18)	66	0.9	1.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.7	1.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	42	0.6	1.0
14	Chronic lower respiratory diseases (J40-J47)	38	0.5	0.9
15	Viral hepatitis (B15-B19)	32	0.4	0.7
...	All other causes (residual)	1,071	14.6	24.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-54 years				
...	All causes	12,145	100.0	349.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,182	18.0	62.8
2	Malignant neoplasms (C00-C97)	2,165	17.8	62.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,109	17.4	60.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,123	9.2	32.3
5	Diabetes mellitus (E10-E14)	606	5.0	17.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	456	3.8	13.1
7	Cerebrovascular diseases (I60-I69)	455	3.7	13.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	246	2.0	7.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	184	1.5	5.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	180	1.5	5.2
11	Septicemia (A40-A41)	166	1.4	4.8
12	Influenza and pneumonia (J09-J18)	143	1.2	4.1
13	Viral hepatitis (B15-B19)	138	1.1	4.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	1.0	3.5
15	Chronic lower respiratory diseases (J40-J47)	73	0.6	2.1
...	All other causes (residual)	1,798	14.8	51.7
Hispanic, male, 55-64 years				
...	All causes	17,974	100.0	787.4
1	Malignant neoplasms (C00-C97)	4,436	24.7	194.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,991	22.2	174.8
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,421	7.9	62.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,366	7.6	59.8
5	Diabetes mellitus (E10-E14)	1,038	5.8	45.5
6	Cerebrovascular diseases (I60-I69)	831	4.6	36.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	338	1.9	14.8
8	Chronic lower respiratory diseases (J40-J47)	283	1.6	12.4
9	Septicemia (A40-A41)	272	1.5	11.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	270	1.5	11.8
11	Viral hepatitis (B15-B19)	253	1.4	11.1
12	Influenza and pneumonia (J09-J18)	251	1.4	11.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	250	1.4	11.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	175	1.0	7.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	0.7	5.8
...	All other causes (residual)	2,666	14.8	116.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-74 years				
...	All causes	19,032	100.0	1,639.9
1	Malignant neoplasms (C00-C97)	5,371	28.2	462.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,515	23.7	389.0
3	Diabetes mellitus (E10-E14)	1,301	6.8	112.1
4	Cerebrovascular diseases (I60-I69)	930	4.9	80.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	794	4.2	68.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	666	3.5	57.4
7	Chronic lower respiratory diseases (J40-J47)	560	2.9	48.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	472	2.5	40.7
9	Influenza and pneumonia (J09-J18)	357	1.9	30.8
10	Septicemia (A40-A41)	302	1.6	26.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	264	1.4	22.7
12	Alzheimer disease (G30)	201	1.1	17.3
13	Parkinson disease (G20-G21)	179	0.9	15.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	133	0.7	11.5
15	Viral hepatitis (B15-B19)	119	0.6	10.3
...	All other causes (residual)	2,868	15.1	247.1
Hispanic, male, 75-84 years				
...	All causes	20,011	100.0	4,086.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,992	24.9	1,019.5
2	Malignant neoplasms (C00-C97)	4,857	24.3	991.9
3	Cerebrovascular diseases (I60-I69)	1,262	6.3	257.7
4	Diabetes mellitus (E10-E14)	1,178	5.9	240.6
5	Chronic lower respiratory diseases (J40-J47)	880	4.4	179.7
6	Alzheimer disease (G30)	705	3.5	144.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	527	2.6	107.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	480	2.4	98.0
9	Influenza and pneumonia (J09-J18)	474	2.4	96.8
10	Parkinson disease (G20-G21)	446	2.2	91.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	320	1.6	65.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	305	1.5	62.3
13	Septicemia (A40-A41)	302	1.5	61.7
14	Pneumonitis due to solids and liquids (J69)	173	0.9	35.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	115	0.6	23.5
...	All other causes (residual)	2,995	15.0	611.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85 years and over				
...	All causes	17,588	100.0	10,147.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,919	28.0	2,837.9
2	Malignant neoplasms (C00-C97)	2,705	15.4	1,560.6
3	Alzheimer disease (G30)	1,344	7.6	775.4
4	Cerebrovascular diseases (I60-I69)	1,229	7.0	709.1
5	Chronic lower respiratory diseases (J40-J47)	860	4.9	496.2
6	Diabetes mellitus (E10-E14)	650	3.7	375.0
7	Influenza and pneumonia (J09-J18)	521	3.0	300.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	417	2.4	240.6
9	Parkinson disease (G20-G21)	397	2.3	229.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	396	2.3	228.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	358	2.0	206.5
12	Septicemia (A40-A41)	275	1.6	158.7
13	Pneumonitis due to solids and liquids (J69)	184	1.0	106.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	119	0.7	68.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	106	0.6	61.2
...	All other causes (residual)	3,108	17.7	1,793.1
Hispanic, female, all ages				
...	All causes	88,670	100.0	304.0
1	Malignant neoplasms (C00-C97)	19,516	22.0	66.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,393	19.6	59.6
3	Cerebrovascular diseases (I60-I69)	5,702	6.4	19.5
4	Alzheimer disease (G30)	5,086	5.7	17.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,302	4.9	14.7
6	Diabetes mellitus (E10-E14)	4,162	4.7	14.3
7	Chronic lower respiratory diseases (J40-J47)	2,840	3.2	9.7
8	Influenza and pneumonia (J09-J18)	1,940	2.2	6.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,940	2.2	6.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,917	2.2	6.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,445	1.6	5.0
12	Septicemia (A40-A41)	1,396	1.6	4.8
13	Certain conditions originating in the perinatal period (P00-P96)	1,053	1.2	3.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	873	1.0	3.0
15	Parkinson disease (G20-G21)	758	0.9	2.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	758	0.9	2.6
...	All other causes (residual)	17,589	19.8	60.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 1-4 years				
...	All causes	350	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	26.0	4.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	16.3	2.8
3	Malignant neoplasms (C00-C97)	42	12.0	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	5.7	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	3.4	*
6	Influenza and pneumonia (J09-J18)	9	2.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	2.3	*
8	Septicemia (A40-A41)	4	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	4	1.1	*
10	Cerebrovascular diseases (I60-I69)	3	0.9	*
11	Anemias (D50-D64)	2	0.6	*
11	Meningitis (G00,G03)	2	0.6	*
13	Meningococcal infection (A39)	1	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (residual)	91	26.0	4.5
Hispanic, female, 5-14 years				
...	All causes	488	100.0	9.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	24.8	2.4
2	Malignant neoplasms (C00-C97)	109	22.3	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	7.0	0.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	6.1	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	4.5	0.4
6	Cerebrovascular diseases (I60-I69)	14	2.9	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.5	*
8	Influenza and pneumonia (J09-J18)	11	2.3	*
9	Septicemia (A40-A41)	10	2.0	*
10	Chronic lower respiratory diseases (J40-J47)	9	1.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.0	*
12	Diseases of appendix (K35-K38)	2	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Anemias (D50-D64)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (residual)	101	20.7	2.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-24 years				
...	All causes	1,467	100.0	31.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	596	40.6	12.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	200	13.6	4.2
3	Malignant neoplasms (C00-C97)	139	9.5	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	132	9.0	2.8
5	Pregnancy, childbirth and the puerperium (O00-O99)	34	2.3	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	2.2	0.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	2.1	0.7
8	Influenza and pneumonia (J09-J18)	16	1.1	*
9	Cerebrovascular diseases (I60-I69)	15	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	14	1.0	*
11	Diabetes mellitus (E10-E14)	10	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.6	*
13	Septicemia (A40-A41)	8	0.5	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	5	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (residual)	222	15.1	4.7
Hispanic, female, 25-34 years				
...	All causes	2,183	100.0	49.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	652	29.9	14.7
2	Malignant neoplasms (C00-C97)	377	17.3	8.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	168	7.7	3.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	150	6.9	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	113	5.2	2.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	80	3.7	1.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	52	2.4	1.2
8	Cerebrovascular diseases (I60-I69)	50	2.3	1.1
9	Diabetes mellitus (E10-E14)	44	2.0	1.0
10	Septicemia (A40-A41)	24	1.1	0.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.0	0.5
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	1.0	0.5
13	Influenza and pneumonia (J09-J18)	20	0.9	0.5
14	Chronic lower respiratory diseases (J40-J47)	16	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.6	*
...	All other causes (residual)	381	17.5	8.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-44 years				
...	All causes	3,380	100.0	81.9
1	Malignant neoplasms (C00-C97)	1,029	30.4	24.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	569	16.8	13.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	309	9.1	7.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	162	4.8	3.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	135	4.0	3.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	117	3.5	2.8
7	Cerebrovascular diseases (I60-I69)	110	3.3	2.7
8	Diabetes mellitus (E10-E14)	93	2.8	2.3
9	Pregnancy, childbirth and the puerperium (O00-O99)	56	1.7	1.4
10	Septicemia (A40-A41)	49	1.4	1.2
11	Influenza and pneumonia (J09-J18)	41	1.2	1.0
12	Chronic lower respiratory diseases (J40-J47)	29	0.9	0.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.9	0.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.8	0.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.7	0.6
...	All other causes (residual)	601	17.8	14.6
Hispanic, female, 45-54 years				
...	All causes	6,264	100.0	183.2
1	Malignant neoplasms (C00-C97)	2,244	35.8	65.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	769	12.3	22.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	534	8.5	15.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	342	5.5	10.0
5	Diabetes mellitus (E10-E14)	328	5.2	9.6
6	Cerebrovascular diseases (I60-I69)	278	4.4	8.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	126	2.0	3.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	113	1.8	3.3
9	Influenza and pneumonia (J09-J18)	110	1.8	3.2
10	Septicemia (A40-A41)	109	1.7	3.2
11	Chronic lower respiratory diseases (J40-J47)	95	1.5	2.8
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	1.1	2.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	0.7	1.3
14	Viral hepatitis (B15-B19)	43	0.7	1.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	39	0.6	1.1
...	All other causes (residual)	1,016	16.2	29.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 55-64 years				
...	All causes	10,559	100.0	436.4
1	Malignant neoplasms (C00-C97)	3,817	36.1	157.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,592	15.1	65.8
3	Diabetes mellitus (E10-E14)	704	6.7	29.1
4	Cerebrovascular diseases (I60-I69)	480	4.5	19.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	476	4.5	19.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	418	4.0	17.3
7	Chronic lower respiratory diseases (J40-J47)	263	2.5	10.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	260	2.5	10.7
9	Septicemia (A40-A41)	193	1.8	8.0
10	Influenza and pneumonia (J09-J18)	184	1.7	7.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	145	1.4	6.0
12	Viral hepatitis (B15-B19)	121	1.1	5.0
13	Alzheimer disease (G30)	61	0.6	2.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	0.5	2.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.4	1.9
...	All other causes (residual)	1,741	16.5	72.0
Hispanic, female, 65-74 years				
...	All causes	14,336	100.0	1,032.1
1	Malignant neoplasms (C00-C97)	4,580	31.9	329.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,632	18.4	189.5
3	Diabetes mellitus (E10-E14)	962	6.7	69.3
4	Cerebrovascular diseases (I60-I69)	784	5.5	56.4
5	Chronic lower respiratory diseases (J40-J47)	474	3.3	34.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	455	3.2	32.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	427	3.0	30.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	323	2.3	23.3
9	Septicemia (A40-A41)	281	2.0	20.2
10	Alzheimer disease (G30)	279	1.9	20.1
11	Influenza and pneumonia (J09-J18)	278	1.9	20.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	246	1.7	17.7
13	Parkinson disease (G20-G21)	91	0.6	6.6
14	Pneumonitis due to solids and liquids (J69)	68	0.5	4.9
15	Viral hepatitis (B15-B19)	56	0.4	4.0
...	All other causes (residual)	2,400	16.7	172.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-84 years				
...	All causes	19,819	100.0	2,877.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,387	22.1	636.8
2	Malignant neoplasms (C00-C97)	4,239	21.4	615.4
3	Cerebrovascular diseases (I60-I69)	1,495	7.5	217.0
4	Alzheimer disease (G30)	1,303	6.6	189.1
5	Diabetes mellitus (E10-E14)	1,134	5.7	164.6
6	Chronic lower respiratory diseases (J40-J47)	806	4.1	117.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	528	2.7	76.6
8	Influenza and pneumonia (J09-J18)	485	2.4	70.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	384	1.9	55.7
10	Septicemia (A40-A41)	362	1.8	52.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	356	1.8	51.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	318	1.6	46.2
13	Parkinson disease (G20-G21)	309	1.6	44.9
14	Pneumonitis due to solids and liquids (J69)	122	0.6	17.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	117	0.6	17.0
...	All other causes (residual)	3,474	17.5	504.3
Hispanic, female, 85 years and over				
...	All causes	27,648	100.0	9,136.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,511	27.2	2,482.0
2	Alzheimer disease (G30)	3,434	12.4	1,134.8
3	Malignant neoplasms (C00-C97)	2,929	10.6	967.9
4	Cerebrovascular diseases (I60-I69)	2,466	8.9	814.9
5	Chronic lower respiratory diseases (J40-J47)	1,131	4.1	373.7
6	Diabetes mellitus (E10-E14)	887	3.2	293.1
7	Influenza and pneumonia (J09-J18)	774	2.8	255.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	633	2.3	209.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	548	2.0	181.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	522	1.9	172.5
11	Septicemia (A40-A41)	341	1.2	112.7
12	Parkinson disease (G20-G21)	328	1.2	108.4
13	Pneumonitis due to solids and liquids (J69)	204	0.7	67.4
14	Nutritional deficiencies (E40-E64)	179	0.6	59.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	119	0.4	39.3
...	All other causes (residual)	5,642	20.4	1,864.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,179,857	100.0	1,083.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	508,485	23.3	252.7
2	Malignant neoplasms (C00-C97)	465,679	21.4	231.4
3	Chronic lower respiratory diseases (J40-J47)	139,833	6.4	69.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127,029	5.8	63.1
5	Cerebrovascular diseases (I60-I69)	110,038	5.0	54.7
6	Alzheimer disease (G30)	101,876	4.7	50.6
7	Diabetes mellitus (E10-E14)	55,116	2.5	27.4
8	Influenza and pneumonia (J09-J18)	43,397	2.0	21.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38,106	1.7	18.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,191	1.6	17.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	30,223	1.4	15.0
12	Septicemia (A40-A41)	30,198	1.4	15.0
13	Parkinson disease (G20-G21)	27,582	1.3	13.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24,465	1.1	12.2
15	Pneumonitis due to solids and liquids (J69)	16,449	0.8	8.2
...	All other causes (residual)	426,190	19.6	211.8
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,858	100.0	22.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	677	36.4	8.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	183	9.8	2.2
3	Malignant neoplasms (C00-C97)	164	8.8	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	113	6.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	3.1	0.7
6	Influenza and pneumonia (J09-J18)	44	2.4	0.5
7	Cerebrovascular diseases (I60-I69)	29	1.6	0.3
8	Septicemia (A40-A41)	24	1.3	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.2	0.3
10	Anemias (D50-D64)	12	0.6	*
11	Acute bronchitis and bronchiolitis (J20-J21)	9	0.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	9	0.5	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	8	0.4	*
15	Meningitis (G00,G03)	7	0.4	*
...	All other causes (residual)	490	26.4	5.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,782	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	819	29.4	3.8
2	Malignant neoplasms (C00-C97)	433	15.6	2.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	328	11.8	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	193	6.9	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	118	4.2	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	3.0	0.4
7	Influenza and pneumonia (J09-J18)	65	2.3	0.3
8	Cerebrovascular diseases (I60-I69)	45	1.6	0.2
9	Chronic lower respiratory diseases (J40-J47)	35	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.0	0.1
11	Septicemia (A40-A41)	19	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.5	*
13	Diabetes mellitus (E10-E14)	11	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	9	0.3	*
15	Anemias (D50-D64)	7	0.3	*
15	Meningitis (G00,G03)	7	0.3	*
15	Pneumonitis due to solids and liquids (J69)	7	0.3	*
...	All other causes (residual)	559	20.1	2.6
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	17,207	100.0	72.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,555	49.7	35.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,074	23.7	17.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	763	4.4	3.2
4	Malignant neoplasms (C00-C97)	686	4.0	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	440	2.6	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	190	1.1	0.8
7	Diabetes mellitus (E10-E14)	129	0.7	0.5
8	Influenza and pneumonia (J09-J18)	100	0.6	0.4
9	Chronic lower respiratory diseases (J40-J47)	71	0.4	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	69	0.4	0.3
11	Cerebrovascular diseases (I60-I69)	66	0.4	0.3
12	Septicemia (A40-A41)	65	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.3	0.2
14	Complications of medical and surgical care (Y40-Y84,Y88)	28	0.2	0.1
15	Legal intervention (Y35,Y89.0)	26	0.2	0.1
...	All other causes (residual)	1,897	11.0	8.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 25-34 years				
...	All causes	36,764	100.0	143.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,243	49.6	71.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,686	15.5	22.2
3	Malignant neoplasms (C00-C97)	1,992	5.4	7.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,962	5.3	7.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,193	3.2	4.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	582	1.6	2.3
7	Diabetes mellitus (E10-E14)	393	1.1	1.5
8	Cerebrovascular diseases (I60-I69)	280	0.8	1.1
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	275	0.7	1.1
10	Pregnancy, childbirth and the puerperium (O00-O99)	244	0.7	1.0
11	Influenza and pneumonia (J09-J18)	224	0.6	0.9
12	Septicemia (A40-A41)	213	0.6	0.8
13	Chronic lower respiratory diseases (J40-J47)	146	0.4	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	91	0.2	0.4
15	Legal intervention (Y35,Y89.0)	90	0.2	0.4
...	All other causes (residual)	5,150	14.0	20.1
Non-Hispanic, white, both sexes, 35-44 years				
...	All causes	49,396	100.0	209.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,129	32.7	68.5
2	Malignant neoplasms (C00-C97)	6,473	13.1	27.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,915	12.0	25.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,764	11.7	24.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,888	3.8	8.0
6	Diabetes mellitus (E10-E14)	1,054	2.1	4.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	993	2.0	4.2
8	Cerebrovascular diseases (I60-I69)	832	1.7	3.5
9	Septicemia (A40-A41)	476	1.0	2.0
10	Influenza and pneumonia (J09-J18)	459	0.9	1.9
10	Chronic lower respiratory diseases (J40-J47)	459	0.9	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	306	0.6	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	301	0.6	1.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	210	0.4	0.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	208	0.4	0.9
...	All other causes (residual)	7,929	16.1	33.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-54 years				
...	All causes	112,765	100.0	416.7
1	Malignant neoplasms (C00-C97)	26,401	23.4	97.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,940	18.6	77.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,356	15.4	64.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,261	6.4	26.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,713	5.1	21.1
6	Diabetes mellitus (E10-E14)	3,624	3.2	13.4
7	Chronic lower respiratory diseases (J40-J47)	3,132	2.8	11.6
8	Cerebrovascular diseases (I60-I69)	2,713	2.4	10.0
9	Septicemia (A40-A41)	1,477	1.3	5.5
10	Influenza and pneumonia (J09-J18)	1,258	1.1	4.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,048	0.9	3.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	994	0.9	3.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	847	0.8	3.1
14	Viral hepatitis (B15-B19)	614	0.5	2.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	559	0.5	2.1
...	All other causes (residual)	18,828	16.7	69.6
Non-Hispanic, white, both sexes, 55-64 years				
...	All causes	264,999	100.0	886.6
1	Malignant neoplasms (C00-C97)	84,084	31.7	281.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,515	20.9	185.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,057	6.4	57.1
4	Chronic lower respiratory diseases (J40-J47)	15,627	5.9	52.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	9,983	3.8	33.4
6	Diabetes mellitus (E10-E14)	9,212	3.5	30.8
7	Cerebrovascular diseases (I60-I69)	7,591	2.9	25.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,162	2.7	24.0
9	Septicemia (A40-A41)	3,879	1.5	13.0
10	Influenza and pneumonia (J09-J18)	3,499	1.3	11.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,039	1.1	10.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,279	0.9	7.6
13	Viral hepatitis (B15-B19)	1,641	0.6	5.5
14	Pneumonitis due to solids and liquids (J69)	1,170	0.4	3.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,162	0.4	3.9
...	All other causes (residual)	42,099	15.9	140.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 65-74 years				
...	All causes	408,427	100.0	1,806.1
1	Malignant neoplasms (C00-C97)	131,875	32.3	583.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	87,648	21.5	387.6
3	Chronic lower respiratory diseases (J40-J47)	34,796	8.5	153.9
4	Cerebrovascular diseases (I60-I69)	15,712	3.8	69.5
5	Diabetes mellitus (E10-E14)	14,230	3.5	62.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,649	2.9	51.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	7,090	1.7	31.4
8	Septicemia (A40-A41)	6,660	1.6	29.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,634	1.6	29.3
10	Influenza and pneumonia (J09-J18)	6,572	1.6	29.1
11	Alzheimer disease (G30)	5,925	1.5	26.2
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,177	1.0	18.5
13	Parkinson disease (G20-G21)	3,892	1.0	17.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,777	0.9	16.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,544	0.6	11.2
...	All other causes (residual)	65,246	16.0	288.5
Non-Hispanic, white, both sexes, 75-84 years				
...	All causes	530,116	100.0	4,612.9
1	Malignant neoplasms (C00-C97)	126,580	23.9	1,101.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	120,435	22.7	1,048.0
3	Chronic lower respiratory diseases (J40-J47)	45,416	8.6	395.2
4	Cerebrovascular diseases (I60-I69)	29,491	5.6	256.6
5	Alzheimer disease (G30)	26,773	5.1	233.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,075	2.7	122.5
7	Diabetes mellitus (E10-E14)	14,034	2.6	122.1
8	Parkinson disease (G20-G21)	11,339	2.1	98.7
9	Influenza and pneumonia (J09-J18)	10,802	2.0	94.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,140	1.9	88.2
11	Septicemia (A40-A41)	8,169	1.5	71.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,759	1.1	50.1
13	Pneumonitis due to solids and liquids (J69)	4,261	0.8	37.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,007	0.8	34.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,586	0.7	31.2
...	All other causes (residual)	95,249	18.0	828.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 85 years and over				
...	All causes	746,124	100.0	14,336.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	215,346	28.9	4,137.6
2	Malignant neoplasms (C00-C97)	86,956	11.7	1,670.8
3	Alzheimer disease (G30)	68,152	9.1	1,309.5
4	Cerebrovascular diseases (I60-I69)	53,230	7.1	1,022.8
5	Chronic lower respiratory diseases (J40-J47)	40,135	5.4	771.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,863	2.9	420.1
7	Influenza and pneumonia (J09-J18)	20,315	2.7	390.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,869	1.9	266.5
9	Diabetes mellitus (E10-E14)	12,425	1.7	238.7
10	Parkinson disease (G20-G21)	11,732	1.6	225.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,551	1.5	221.9
12	Septicemia (A40-A41)	9,163	1.2	176.1
13	Pneumonitis due to solids and liquids (J69)	8,060	1.1	154.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,680	0.6	89.9
15	Nutritional deficiencies (E40-E64)	3,523	0.5	67.7
...	All other causes (residual)	165,124	22.1	3,172.7
Non-Hispanic, white, male, all ages				
...	All causes	1,102,838	100.0	1,111.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	272,428	24.7	274.5
2	Malignant neoplasms (C00-C97)	247,022	22.4	248.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79,280	7.2	79.9
4	Chronic lower respiratory diseases (J40-J47)	64,673	5.9	65.2
5	Cerebrovascular diseases (I60-I69)	45,078	4.1	45.4
6	Alzheimer disease (G30)	31,577	2.9	31.8
7	Diabetes mellitus (E10-E14)	31,343	2.8	31.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29,708	2.7	29.9
9	Influenza and pneumonia (J09-J18)	20,515	1.9	20.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	18,866	1.7	19.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,187	1.6	18.3
12	Parkinson disease (G20-G21)	16,841	1.5	17.0
13	Septicemia (A40-A41)	14,419	1.3	14.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,696	1.0	10.8
15	Pneumonitis due to solids and liquids (J69)	9,352	0.8	9.4
...	All other causes (residual)	192,853	17.5	194.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,107	100.0	26.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	425	38.4	10.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	9.1	2.4
3	Malignant neoplasms (C00-C97)	93	8.4	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	5.8	1.5
5	Influenza and pneumonia (J09-J18)	30	2.7	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	2.6	0.7
7	Cerebrovascular diseases (I60-I69)	18	1.6	*
8	Septicemia (A40-A41)	12	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
11	Meningitis (G00,G03)	6	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
13	Anemias (D50-D64)	4	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	4	0.4	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
...	All other causes (residual)	295	26.6	6.9
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,628	100.0	14.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	514	31.6	4.6
2	Malignant neoplasms (C00-C97)	235	14.4	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	227	13.9	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	103	6.3	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	3.8	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	2.5	0.4
7	Cerebrovascular diseases (I60-I69)	27	1.7	0.2
8	Influenza and pneumonia (J09-J18)	25	1.5	0.2
9	Chronic lower respiratory diseases (J40-J47)	19	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.9	*
11	Septicemia (A40-A41)	13	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.5	*
13	Diabetes mellitus (E10-E14)	6	0.4	*
14	Anemias (D50-D64)	5	0.3	*
14	Meningitis (G00,G03)	5	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
...	All other causes (residual)	318	19.5	2.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 15-24 years				
...	All causes	12,285	100.0	100.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,120	49.8	50.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,328	27.1	27.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	538	4.4	4.4
4	Malignant neoplasms (C00-C97)	413	3.4	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	292	2.4	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	110	0.9	0.9
7	Diabetes mellitus (E10-E14)	74	0.6	0.6
8	Influenza and pneumonia (J09-J18)	56	0.5	0.5
9	Chronic lower respiratory diseases (J40-J47)	44	0.4	0.4
10	Cerebrovascular diseases (I60-I69)	32	0.3	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.2	0.2
12	Septicemia (A40-A41)	27	0.2	0.2
13	Legal intervention (Y35,Y89.0)	22	0.2	0.2
14	Aortic aneurysm and dissection (I71)	18	0.1	*
14	Pneumonitis due to solids and liquids (J69)	18	0.1	*
...	All other causes (residual)	1,163	9.5	9.5
Non-Hispanic, white, male, 25-34 years				
...	All causes	25,360	100.0	194.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,281	52.4	102.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,527	17.9	34.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,273	5.0	9.8
4	Malignant neoplasms (C00-C97)	949	3.7	7.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	849	3.3	6.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	340	1.3	2.6
7	Diabetes mellitus (E10-E14)	234	0.9	1.8
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	177	0.7	1.4
9	Cerebrovascular diseases (I60-I69)	171	0.7	1.3
10	Influenza and pneumonia (J09-J18)	117	0.5	0.9
11	Septicemia (A40-A41)	102	0.4	0.8
12	Legal intervention (Y35,Y89.0)	85	0.3	0.7
13	Chronic lower respiratory diseases (J40-J47)	75	0.3	0.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	55	0.2	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.2	0.4
...	All other causes (residual)	3,072	12.1	23.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 35-44 years				
...	All causes	31,274	100.0	264.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,125	35.6	94.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,385	14.0	37.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,076	13.0	34.4
4	Malignant neoplasms (C00-C97)	2,925	9.4	24.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,124	3.6	9.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	670	2.1	5.7
7	Diabetes mellitus (E10-E14)	649	2.1	5.5
8	Cerebrovascular diseases (I60-I69)	447	1.4	3.8
9	Septicemia (A40-A41)	253	0.8	2.1
10	Influenza and pneumonia (J09-J18)	244	0.8	2.1
11	Chronic lower respiratory diseases (J40-J47)	195	0.6	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	180	0.6	1.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	167	0.5	1.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	159	0.5	1.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	147	0.5	1.2
...	All other causes (residual)	4,528	14.5	38.2
Non-Hispanic, white, male, 45-54 years				
...	All causes	68,781	100.0	510.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,885	21.6	110.4
2	Malignant neoplasms (C00-C97)	12,975	18.9	96.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,642	16.9	86.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,331	7.8	39.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,594	5.2	26.7
6	Diabetes mellitus (E10-E14)	2,311	3.4	17.1
7	Cerebrovascular diseases (I60-I69)	1,553	2.3	11.5
8	Chronic lower respiratory diseases (J40-J47)	1,446	2.1	10.7
9	Septicemia (A40-A41)	796	1.2	5.9
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	671	1.0	5.0
11	Influenza and pneumonia (J09-J18)	656	1.0	4.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	611	0.9	4.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	552	0.8	4.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	461	0.7	3.4
15	Viral hepatitis (B15-B19)	387	0.6	2.9
...	All other causes (residual)	10,910	15.9	80.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 55-64 years				
...	All causes	161,298	100.0	1,105.5
1	Malignant neoplasms (C00-C97)	46,162	28.6	316.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,100	24.2	268.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,533	7.2	79.0
4	Chronic lower respiratory diseases (J40-J47)	7,896	4.9	54.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,581	4.1	45.1
6	Diabetes mellitus (E10-E14)	5,911	3.7	40.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,403	3.3	37.0
8	Cerebrovascular diseases (I60-I69)	4,320	2.7	29.6
9	Septicemia (A40-A41)	2,102	1.3	14.4
10	Influenza and pneumonia (J09-J18)	1,964	1.2	13.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,772	1.1	12.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,456	0.9	10.0
13	Viral hepatitis (B15-B19)	1,132	0.7	7.8
14	Pneumonitis due to solids and liquids (J69)	743	0.5	5.1
15	Aortic aneurysm and dissection (I71)	705	0.4	4.8
...	All other causes (residual)	24,518	15.2	168.0
Non-Hispanic, white, male, 65-74 years				
...	All causes	234,580	100.0	2,182.5
1	Malignant neoplasms (C00-C97)	73,185	31.2	680.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	57,064	24.3	530.9
3	Chronic lower respiratory diseases (J40-J47)	17,454	7.4	162.4
4	Diabetes mellitus (E10-E14)	8,623	3.7	80.2
5	Cerebrovascular diseases (I60-I69)	8,325	3.5	77.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,478	3.2	69.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,560	1.9	42.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,720	1.6	34.6
9	Influenza and pneumonia (J09-J18)	3,539	1.5	32.9
10	Septicemia (A40-A41)	3,379	1.4	31.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,297	1.4	30.7
12	Parkinson disease (G20-G21)	2,589	1.1	24.1
13	Alzheimer disease (G30)	2,526	1.1	23.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,157	0.9	20.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,520	0.6	14.1
...	All other causes (residual)	35,164	15.0	327.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 75-84 years				
...	All causes	274,094	100.0	5,378.9
1	Malignant neoplasms (C00-C97)	68,102	24.8	1,336.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	67,956	24.8	1,333.6
3	Chronic lower respiratory diseases (J40-J47)	21,166	7.7	415.4
4	Cerebrovascular diseases (I60-I69)	13,461	4.9	264.2
5	Alzheimer disease (G30)	10,128	3.7	198.8
6	Diabetes mellitus (E10-E14)	7,960	2.9	156.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,781	2.8	152.7
8	Parkinson disease (G20-G21)	7,369	2.7	144.6
9	Influenza and pneumonia (J09-J18)	5,549	2.0	108.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,391	2.0	105.8
11	Septicemia (A40-A41)	3,930	1.4	77.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,706	1.0	53.1
13	Pneumonitis due to solids and liquids (J69)	2,548	0.9	50.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,250	0.8	44.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,122	0.8	41.6
...	All other causes (residual)	45,675	16.7	896.3
Non-Hispanic, white, male, 85 years and over				
...	All causes	287,103	100.0	15,521.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	87,636	30.5	4,737.9
2	Malignant neoplasms (C00-C97)	41,963	14.6	2,268.7
3	Alzheimer disease (G30)	18,470	6.4	998.6
4	Cerebrovascular diseases (I60-I69)	16,695	5.8	902.6
5	Chronic lower respiratory diseases (J40-J47)	16,371	5.7	885.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,011	3.1	487.2
7	Influenza and pneumonia (J09-J18)	8,299	2.9	448.7
8	Parkinson disease (G20-G21)	6,465	2.3	349.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,428	2.2	347.5
10	Diabetes mellitus (E10-E14)	5,573	1.9	301.3
11	Pneumonitis due to solids and liquids (J69)	4,227	1.5	228.5
12	Septicemia (A40-A41)	3,775	1.3	204.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,648	1.3	197.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,238	0.8	121.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,087	0.4	58.8
...	All other causes (residual)	55,217	19.2	2,985.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, all ages				
...	All causes	1,077,019	100.0	1,055.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	236,057	21.9	231.4
2	Malignant neoplasms (C00-C97)	218,657	20.3	214.3
3	Chronic lower respiratory diseases (J40-J47)	75,160	7.0	73.7
4	Alzheimer disease (G30)	70,299	6.5	68.9
5	Cerebrovascular diseases (I60-I69)	64,960	6.0	63.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47,749	4.4	46.8
7	Diabetes mellitus (E10-E14)	23,773	2.2	23.3
8	Influenza and pneumonia (J09-J18)	22,882	2.1	22.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,004	1.6	16.7
10	Septicemia (A40-A41)	15,779	1.5	15.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,769	1.3	13.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	11,357	1.1	11.1
13	Parkinson disease (G20-G21)	10,741	1.0	10.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,398	0.8	8.2
15	Pneumonitis due to solids and liquids (J69)	7,097	0.7	7.0
...	All other causes (residual)	233,337	21.7	228.7
Non-Hispanic, white, female, 1-4 years				
...	All causes	751	100.0	18.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	252	33.6	6.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	10.9	2.0
3	Malignant neoplasms (C00-C97)	71	9.5	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	6.5	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.9	0.7
6	Influenza and pneumonia (J09-J18)	14	1.9	*
7	Septicemia (A40-A41)	12	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.5	*
8	Cerebrovascular diseases (I60-I69)	11	1.5	*
10	Anemias (D50-D64)	8	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.7	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
14	Hernia (K40-K46)	3	0.4	*
15	Meningococcal infection (A39)	2	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.3	*
15	Pneumonitis due to solids and liquids (J69)	2	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	184	24.5	4.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,154	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	26.4	2.9
2	Malignant neoplasms (C00-C97)	198	17.2	1.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	8.8	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	7.8	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	4.9	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	3.7	0.4
7	Influenza and pneumonia (J09-J18)	40	3.5	0.4
8	Cerebrovascular diseases (I60-I69)	18	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	16	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.1	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
12	Septicemia (A40-A41)	6	0.5	*
13	Diabetes mellitus (E10-E14)	5	0.4	*
14	Pneumonitis due to solids and liquids (J69)	4	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	4	0.3	*
...	All other causes (residual)	248	21.5	2.3
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,922	100.0	42.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,435	49.5	21.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	746	15.2	6.4
3	Malignant neoplasms (C00-C97)	273	5.5	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	225	4.6	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	148	3.0	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	1.6	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	69	1.4	0.6
8	Diabetes mellitus (E10-E14)	55	1.1	0.5
9	Influenza and pneumonia (J09-J18)	44	0.9	0.4
10	Septicemia (A40-A41)	38	0.8	0.3
11	Cerebrovascular diseases (I60-I69)	34	0.7	0.3
12	Chronic lower respiratory diseases (J40-J47)	27	0.5	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.3	*
...	All other causes (residual)	704	14.3	6.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 25-34 years				
...	All causes	11,404	100.0	90.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,962	43.5	39.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,159	10.2	9.2
3	Malignant neoplasms (C00-C97)	1,043	9.1	8.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	689	6.0	5.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	344	3.0	2.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	244	2.1	1.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	242	2.1	1.9
8	Diabetes mellitus (E10-E14)	159	1.4	1.3
9	Septicemia (A40-A41)	111	1.0	0.9
10	Cerebrovascular diseases (I60-I69)	109	1.0	0.9
11	Influenza and pneumonia (J09-J18)	107	0.9	0.8
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	98	0.9	0.8
13	Chronic lower respiratory diseases (J40-J47)	71	0.6	0.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.4	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	0.3	0.3
...	All other causes (residual)	1,988	17.4	15.7
Non-Hispanic, white, female, 35-44 years				
...	All causes	18,122	100.0	154.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,004	27.6	42.8
2	Malignant neoplasms (C00-C97)	3,548	19.6	30.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,839	10.1	15.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,379	7.6	11.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	764	4.2	6.5
6	Diabetes mellitus (E10-E14)	405	2.2	3.5
7	Cerebrovascular diseases (I60-I69)	385	2.1	3.3
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	323	1.8	2.8
9	Chronic lower respiratory diseases (J40-J47)	264	1.5	2.3
10	Septicemia (A40-A41)	223	1.2	1.9
11	Influenza and pneumonia (J09-J18)	215	1.2	1.8
12	Pregnancy, childbirth and the puerperium (O00-O99)	152	0.8	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	134	0.7	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	0.7	1.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	61	0.3	0.5
...	All other causes (residual)	3,300	18.2	28.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-54 years				
...	All causes	43,984	100.0	323.9
1	Malignant neoplasms (C00-C97)	13,426	30.5	98.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,055	13.8	44.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,714	13.0	42.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,119	4.8	15.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,930	4.4	14.2
6	Chronic lower respiratory diseases (J40-J47)	1,686	3.8	12.4
7	Diabetes mellitus (E10-E14)	1,313	3.0	9.7
8	Cerebrovascular diseases (I60-I69)	1,160	2.6	8.5
9	Septicemia (A40-A41)	681	1.5	5.0
10	Influenza and pneumonia (J09-J18)	602	1.4	4.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	437	1.0	3.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	323	0.7	2.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	295	0.7	2.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	262	0.6	1.9
15	Viral hepatitis (B15-B19)	227	0.5	1.7
...	All other causes (residual)	7,754	17.6	57.1
Non-Hispanic, white, female, 55-64 years				
...	All causes	103,701	100.0	677.9
1	Malignant neoplasms (C00-C97)	37,922	36.6	247.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,415	15.8	107.3
3	Chronic lower respiratory diseases (J40-J47)	7,731	7.5	50.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,524	5.3	36.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,402	3.3	22.2
6	Diabetes mellitus (E10-E14)	3,301	3.2	21.6
7	Cerebrovascular diseases (I60-I69)	3,271	3.2	21.4
8	Septicemia (A40-A41)	1,777	1.7	11.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,759	1.7	11.5
10	Influenza and pneumonia (J09-J18)	1,535	1.5	10.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,267	1.2	8.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	823	0.8	5.4
13	Alzheimer disease (G30)	525	0.5	3.4
14	Viral hepatitis (B15-B19)	509	0.5	3.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	501	0.5	3.3
...	All other causes (residual)	17,439	16.8	114.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 65-74 years				
...	All causes	173,847	100.0	1,465.2
1	Malignant neoplasms (C00-C97)	58,690	33.8	494.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,584	17.6	257.8
3	Chronic lower respiratory diseases (J40-J47)	17,342	10.0	146.2
4	Cerebrovascular diseases (I60-I69)	7,387	4.2	62.3
5	Diabetes mellitus (E10-E14)	5,607	3.2	47.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,171	2.4	35.2
7	Alzheimer disease (G30)	3,399	2.0	28.6
8	Septicemia (A40-A41)	3,281	1.9	27.7
9	Influenza and pneumonia (J09-J18)	3,033	1.7	25.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,914	1.7	24.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,530	1.5	21.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,620	0.9	13.7
13	Parkinson disease (G20-G21)	1,303	0.7	11.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,024	0.6	8.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	880	0.5	7.4
...	All other causes (residual)	30,082	17.3	253.5
Non-Hispanic, white, female, 75-84 years				
...	All causes	256,022	100.0	4,002.6
1	Malignant neoplasms (C00-C97)	58,478	22.8	914.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,479	20.5	820.4
3	Chronic lower respiratory diseases (J40-J47)	24,250	9.5	379.1
4	Alzheimer disease (G30)	16,645	6.5	260.2
5	Cerebrovascular diseases (I60-I69)	16,030	6.3	250.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,294	2.5	98.4
7	Diabetes mellitus (E10-E14)	6,074	2.4	95.0
8	Influenza and pneumonia (J09-J18)	5,253	2.1	82.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,749	1.9	74.2
10	Septicemia (A40-A41)	4,239	1.7	66.3
11	Parkinson disease (G20-G21)	3,970	1.6	62.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,053	1.2	47.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,757	0.7	27.5
14	Pneumonitis due to solids and liquids (J69)	1,713	0.7	26.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,565	0.6	24.5
...	All other causes (residual)	49,473	19.3	773.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 85 years and over				
...	All causes	459,021	100.0	13,682.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	127,710	27.8	3,806.7
2	Alzheimer disease (G30)	49,682	10.8	1,480.9
3	Malignant neoplasms (C00-C97)	44,993	9.8	1,341.1
4	Cerebrovascular diseases (I60-I69)	36,535	8.0	1,089.0
5	Chronic lower respiratory diseases (J40-J47)	23,764	5.2	708.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,852	2.8	383.1
7	Influenza and pneumonia (J09-J18)	12,016	2.6	358.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,903	1.7	235.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,441	1.6	221.8
10	Diabetes mellitus (E10-E14)	6,852	1.5	204.2
11	Septicemia (A40-A41)	5,388	1.2	160.6
12	Parkinson disease (G20-G21)	5,267	1.1	157.0
13	Pneumonitis due to solids and liquids (J69)	3,833	0.8	114.3
14	Nutritional deficiencies (E40-E64)	2,460	0.5	73.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,442	0.5	72.8
...	All other causes (residual)	109,883	23.9	3,275.3
Non-Hispanic, black, both sexes, all ages				
...	All causes	335,667	100.0	787.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	78,161	23.3	183.4
2	Malignant neoplasms (C00-C97)	69,872	20.8	163.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,869	5.9	46.6
4	Cerebrovascular diseases (I60-I69)	19,088	5.7	44.8
5	Diabetes mellitus (E10-E14)	14,798	4.4	34.7
6	Chronic lower respiratory diseases (J40-J47)	11,217	3.3	26.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9,908	3.0	23.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,542	2.8	22.4
9	Alzheimer disease (G30)	8,991	2.7	21.1
10	Septicemia (A40-A41)	6,568	2.0	15.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,273	1.9	14.7
12	Influenza and pneumonia (J09-J18)	5,556	1.7	13.0
13	Certain conditions originating in the perinatal period (P00-P96)	3,637	1.1	8.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	3,290	1.0	7.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2,965	0.9	7.0
...	All other causes (residual)	65,932	19.6	154.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	985	100.0	40.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	295	29.9	12.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	12.7	5.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	10.3	4.1
4	Malignant neoplasms (C00-C97)	42	4.3	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.8	1.5
6	Influenza and pneumonia (J09-J18)	25	2.5	1.0
7	Cerebrovascular diseases (I60-I69)	21	2.1	0.9
8	Chronic lower respiratory diseases (J40-J47)	18	1.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	18	1.8	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	10	1.0	*
11	Septicemia (A40-A41)	9	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
13	Anemias (D50-D64)	6	0.6	*
13	Meningitis (G00,G03)	6	0.6	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.6	*
...	All other causes (residual)	259	26.3	10.6
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,296	100.0	20.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	377	29.1	6.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	142	11.0	2.3
3	Malignant neoplasms (C00-C97)	136	10.5	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	6.2	1.3
5	Chronic lower respiratory diseases (J40-J47)	77	5.9	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	5.1	1.1
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	3.6	0.8
8	Cerebrovascular diseases (I60-I69)	25	1.9	0.4
9	Influenza and pneumonia (J09-J18)	23	1.8	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.4	*
11	Diabetes mellitus (E10-E14)	16	1.2	*
12	Septicemia (A40-A41)	14	1.1	*
13	Anemias (D50-D64)	12	0.9	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (residual)	253	19.5	4.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	7,364	100.0	111.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,055	41.5	46.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,830	24.9	27.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	710	9.6	10.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	294	4.0	4.4
5	Malignant neoplasms (C00-C97)	219	3.0	3.3
6	Diabetes mellitus (E10-E14)	83	1.1	1.3
7	Chronic lower respiratory diseases (J40-J47)	82	1.1	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	78	1.1	1.2
9	Pregnancy, childbirth and the puerperium (O00-O99)	58	0.8	0.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	47	0.6	0.7
11	Anemias (D50-D64)	44	0.6	0.7
12	Influenza and pneumonia (J09-J18)	41	0.6	0.6
13	Cerebrovascular diseases (I60-I69)	38	0.5	0.6
14	Legal intervention (Y35,Y89.0)	37	0.5	0.6
15	Septicemia (A40-A41)	29	0.4	0.4
...	All other causes (residual)	719	9.8	10.9
Non-Hispanic, black, both sexes, 25-34 years				
...	All causes	12,084	100.0	186.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,196	26.4	49.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,070	25.4	47.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,058	8.8	16.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	793	6.6	12.2
5	Malignant neoplasms (C00-C97)	653	5.4	10.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	335	2.8	5.2
7	Diabetes mellitus (E10-E14)	281	2.3	4.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	161	1.3	2.5
9	Cerebrovascular diseases (I60-I69)	142	1.2	2.2
10	Chronic lower respiratory diseases (J40-J47)	128	1.1	2.0
11	Anemias (D50-D64)	126	1.0	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	102	0.8	1.6
13	Septicemia (A40-A41)	98	0.8	1.5
14	Influenza and pneumonia (J09-J18)	96	0.8	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	0.7	1.2
...	All other causes (residual)	1,766	14.6	27.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 35-44 years				
...	All causes	15,726	100.0	293.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,951	18.8	55.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,839	18.1	53.0
3	Malignant neoplasms (C00-C97)	1,926	12.2	35.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,662	10.6	31.0
5	Diabetes mellitus (E10-E14)	646	4.1	12.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	515	3.3	9.6
7	Cerebrovascular diseases (I60-I69)	475	3.0	8.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	449	2.9	8.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	266	1.7	5.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	221	1.4	4.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	219	1.4	4.1
12	Septicemia (A40-A41)	215	1.4	4.0
13	Influenza and pneumonia (J09-J18)	170	1.1	3.2
14	Chronic lower respiratory diseases (J40-J47)	162	1.0	3.0
15	Anemias (D50-D64)	128	0.8	2.4
...	All other causes (residual)	2,882	18.3	53.8
Non-Hispanic, black, both sexes, 45-54 years				
...	All causes	31,226	100.0	583.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,452	23.9	139.2
2	Malignant neoplasms (C00-C97)	6,260	20.0	116.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,557	11.4	66.4
4	Diabetes mellitus (E10-E14)	1,496	4.8	27.9
5	Cerebrovascular diseases (I60-I69)	1,402	4.5	26.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	869	2.8	16.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	773	2.5	14.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	749	2.4	14.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	659	2.1	12.3
10	Septicemia (A40-A41)	594	1.9	11.1
11	Chronic lower respiratory diseases (J40-J47)	577	1.8	10.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	549	1.8	10.3
13	Influenza and pneumonia (J09-J18)	404	1.3	7.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	398	1.3	7.4
15	Aortic aneurysm and dissection (I71)	162	0.5	3.0
...	All other causes (residual)	5,325	17.1	99.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 55-64 years				
...	All causes	64,498	100.0	1,316.1
1	Malignant neoplasms (C00-C97)	17,750	27.5	362.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,128	25.0	329.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,719	5.8	75.9
4	Diabetes mellitus (E10-E14)	3,193	5.0	65.2
5	Cerebrovascular diseases (I60-I69)	3,162	4.9	64.5
6	Chronic lower respiratory diseases (J40-J47)	2,070	3.2	42.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,798	2.8	36.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,344	2.1	27.4
9	Septicemia (A40-A41)	1,301	2.0	26.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,140	1.8	23.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	911	1.4	18.6
12	Influenza and pneumonia (J09-J18)	888	1.4	18.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	511	0.8	10.4
14	Viral hepatitis (B15-B19)	380	0.6	7.8
15	Pneumonitis due to solids and liquids (J69)	279	0.4	5.7
...	All other causes (residual)	9,924	15.4	202.5
Non-Hispanic, black, both sexes, 65-74 years				
...	All causes	71,021	100.0	2,460.9
1	Malignant neoplasms (C00-C97)	20,610	29.0	714.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,693	24.9	613.1
3	Cerebrovascular diseases (I60-I69)	4,169	5.9	144.5
4	Diabetes mellitus (E10-E14)	3,794	5.3	131.5
5	Chronic lower respiratory diseases (J40-J47)	3,081	4.3	106.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,353	3.3	81.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,768	2.5	61.3
8	Septicemia (A40-A41)	1,643	2.3	56.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,304	1.8	45.2
10	Influenza and pneumonia (J09-J18)	1,169	1.6	40.5
11	Alzheimer disease (G30)	738	1.0	25.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	727	1.0	25.2
13	Pneumonitis due to solids and liquids (J69)	418	0.6	14.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	344	0.5	11.9
15	Parkinson disease (G20-G21)	324	0.5	11.2
...	All other causes (residual)	10,886	15.3	377.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 75-84 years				
...	All causes	65,076	100.0	5,093.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,234	24.9	1,270.5
2	Malignant neoplasms (C00-C97)	14,767	22.7	1,155.7
3	Cerebrovascular diseases (I60-I69)	4,710	7.2	368.6
4	Diabetes mellitus (E10-E14)	3,181	4.9	249.0
5	Chronic lower respiratory diseases (J40-J47)	2,912	4.5	227.9
6	Alzheimer disease (G30)	2,768	4.3	216.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,380	3.7	186.3
8	Septicemia (A40-A41)	1,485	2.3	116.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,341	2.1	105.0
10	Influenza and pneumonia (J09-J18)	1,319	2.0	103.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,015	1.6	79.4
12	Parkinson disease (G20-G21)	645	1.0	50.5
13	Pneumonitis due to solids and liquids (J69)	481	0.7	37.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	346	0.5	27.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	224	0.3	17.5
...	All other causes (residual)	11,268	17.3	881.9
Non-Hispanic, black, both sexes, 85 years and over				
...	All causes	59,645	100.0	12,101.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,281	27.3	3,303.4
2	Malignant neoplasms (C00-C97)	7,500	12.6	1,521.7
3	Alzheimer disease (G30)	5,358	9.0	1,087.1
4	Cerebrovascular diseases (I60-I69)	4,907	8.2	995.6
5	Diabetes mellitus (E10-E14)	2,105	3.5	427.1
5	Chronic lower respiratory diseases (J40-J47)	2,105	3.5	427.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,851	3.1	375.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,657	2.8	336.2
9	Influenza and pneumonia (J09-J18)	1,366	2.3	277.2
10	Septicemia (A40-A41)	1,129	1.9	229.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	831	1.4	168.6
12	Pneumonitis due to solids and liquids (J69)	581	1.0	117.9
13	Parkinson disease (G20-G21)	479	0.8	97.2
14	Nutritional deficiencies (E40-E64)	383	0.6	77.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	264	0.4	53.6
...	All other causes (residual)	12,848	21.5	2,606.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, all ages				
...	All causes	174,403	100.0	854.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,411	23.7	202.8
2	Malignant neoplasms (C00-C97)	35,261	20.2	172.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,844	7.9	67.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,643	5.0	42.3
5	Cerebrovascular diseases (I60-I69)	8,566	4.9	42.0
6	Diabetes mellitus (E10-E14)	7,494	4.3	36.7
7	Chronic lower respiratory diseases (J40-J47)	5,605	3.2	27.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,555	2.6	22.3
9	Septicemia (A40-A41)	3,052	1.7	14.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,874	1.6	14.1
11	Influenza and pneumonia (J09-J18)	2,682	1.5	13.1
12	Alzheimer disease (G30)	2,476	1.4	12.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,324	1.3	11.4
14	Certain conditions originating in the perinatal period (P00-P96)	2,080	1.2	10.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,064	1.2	10.1
...	All other causes (residual)	31,472	18.0	154.2
Non-Hispanic, black, male, 1-4 years				
...	All causes	535	100.0	42.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	30.8	13.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	12.9	5.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	9.3	4.0
4	Malignant neoplasms (C00-C97)	23	4.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	3.9	1.7
6	Influenza and pneumonia (J09-J18)	13	2.4	*
7	Cerebrovascular diseases (I60-I69)	11	2.1	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.9	*
9	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.1	*
11	Septicemia (A40-A41)	4	0.7	*
11	Meningitis (G00,G03)	4	0.7	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
14	Anemias (D50-D64)	3	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
...	All other causes (residual)	144	26.9	11.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 5-14 years				
...	All causes	784	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	247	31.5	7.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	11.1	2.8
3	Malignant neoplasms (C00-C97)	73	9.3	2.3
4	Chronic lower respiratory diseases (J40-J47)	54	6.9	1.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	5.9	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	5.7	1.4
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.9	0.7
8	Cerebrovascular diseases (I60-I69)	15	1.9	*
9	Influenza and pneumonia (J09-J18)	11	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.1	*
10	Diabetes mellitus (E10-E14)	9	1.1	*
12	Septicemia (A40-A41)	8	1.0	*
13	Anemias (D50-D64)	7	0.9	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
15	Pneumonitis due to solids and liquids (J69)	2	0.3	*
...	All other causes (residual)	145	18.5	4.6
Non-Hispanic, black, male, 15-24 years				
...	All causes	5,711	100.0	170.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,761	48.3	82.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,328	23.3	39.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	565	9.9	16.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	199	3.5	5.9
5	Malignant neoplasms (C00-C97)	127	2.2	3.8
6	Chronic lower respiratory diseases (J40-J47)	52	0.9	1.5
7	Diabetes mellitus (E10-E14)	47	0.8	1.4
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	0.8	1.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	37	0.6	1.1
10	Legal intervention (Y35,Y89.0)	36	0.6	1.1
11	Influenza and pneumonia (J09-J18)	25	0.4	0.7
12	Cerebrovascular diseases (I60-I69)	19	0.3	*
13	Anemias (D50-D64)	17	0.3	*
14	Septicemia (A40-A41)	16	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
...	All other causes (residual)	426	7.5	12.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 25-34 years				
...	All causes	8,552	100.0	267.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,865	33.5	89.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,271	26.6	71.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	673	7.9	21.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	652	7.6	20.4
5	Malignant neoplasms (C00-C97)	294	3.4	9.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	231	2.7	7.2
7	Diabetes mellitus (E10-E14)	141	1.6	4.4
8	Chronic lower respiratory diseases (J40-J47)	80	0.9	2.5
9	Anemias (D50-D64)	70	0.8	2.2
10	Cerebrovascular diseases (I60-I69)	67	0.8	2.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.6	1.7
12	Septicemia (A40-A41)	49	0.6	1.5
13	Influenza and pneumonia (J09-J18)	46	0.5	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	0.5	1.4
15	Legal intervention (Y35,Y89.0)	42	0.5	1.3
...	All other causes (residual)	973	11.4	30.4
Non-Hispanic, black, male, 35-44 years				
...	All causes	9,721	100.0	383.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,188	22.5	86.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,787	18.4	70.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,459	15.0	57.6
4	Malignant neoplasms (C00-C97)	751	7.7	29.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	391	4.0	15.4
6	Diabetes mellitus (E10-E14)	380	3.9	15.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	267	2.7	10.5
8	Cerebrovascular diseases (I60-I69)	253	2.6	10.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	141	1.5	5.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	1.4	5.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	114	1.2	4.5
12	Septicemia (A40-A41)	113	1.2	4.5
13	Influenza and pneumonia (J09-J18)	96	1.0	3.8
14	Aortic aneurysm and dissection (I71)	84	0.9	3.3
15	Chronic lower respiratory diseases (J40-J47)	82	0.8	3.2
...	All other causes (residual)	1,476	15.2	58.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-54 years				
...	All causes	18,181	100.0	724.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,760	26.2	189.7
2	Malignant neoplasms (C00-C97)	2,767	15.2	110.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,499	13.7	99.6
4	Diabetes mellitus (E10-E14)	881	4.8	35.1
5	Cerebrovascular diseases (I60-I69)	801	4.4	31.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	751	4.1	29.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	475	2.6	18.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	426	2.3	17.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	378	2.1	15.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	328	1.8	13.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	317	1.7	12.6
12	Septicemia (A40-A41)	307	1.7	12.2
13	Chronic lower respiratory diseases (J40-J47)	260	1.4	10.4
14	Influenza and pneumonia (J09-J18)	218	1.2	8.7
15	Aortic aneurysm and dissection (I71)	115	0.6	4.6
...	All other causes (residual)	2,898	15.9	115.5
Non-Hispanic, black, male, 55-64 years				
...	All causes	37,552	100.0	1,676.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,193	27.1	455.1
2	Malignant neoplasms (C00-C97)	9,321	24.8	416.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,741	7.3	122.4
4	Diabetes mellitus (E10-E14)	1,833	4.9	81.8
5	Cerebrovascular diseases (I60-I69)	1,807	4.8	80.7
6	Chronic lower respiratory diseases (J40-J47)	1,016	2.7	45.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	947	2.5	42.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	889	2.4	39.7
9	Septicemia (A40-A41)	678	1.8	30.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	654	1.7	29.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	617	1.6	27.5
12	Influenza and pneumonia (J09-J18)	483	1.3	21.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	417	1.1	18.6
14	Viral hepatitis (B15-B19)	271	0.7	12.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	0.5	8.3
...	All other causes (residual)	5,498	14.6	245.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 65-74 years				
...	All causes	39,396	100.0	3,179.0
1	Malignant neoplasms (C00-C97)	11,224	28.5	905.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,426	26.5	841.3
3	Cerebrovascular diseases (I60-I69)	2,195	5.6	177.1
4	Diabetes mellitus (E10-E14)	2,066	5.2	166.7
5	Chronic lower respiratory diseases (J40-J47)	1,679	4.3	135.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,256	3.2	101.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,204	3.1	97.2
8	Septicemia (A40-A41)	805	2.0	65.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	672	1.7	54.2
10	Influenza and pneumonia (J09-J18)	645	1.6	52.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	503	1.3	40.6
12	Alzheimer disease (G30)	304	0.8	24.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	257	0.7	20.7
14	Pneumonitis due to solids and liquids (J69)	247	0.6	19.9
15	Parkinson disease (G20-G21)	204	0.5	16.5
...	All other causes (residual)	5,709	14.5	460.7
Non-Hispanic, black, male, 75-84 years				
...	All causes	30,739	100.0	6,248.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,927	25.8	1,611.3
2	Malignant neoplasms (C00-C97)	7,447	24.2	1,513.8
3	Cerebrovascular diseases (I60-I69)	2,024	6.6	411.4
4	Chronic lower respiratory diseases (J40-J47)	1,474	4.8	299.6
5	Diabetes mellitus (E10-E14)	1,456	4.7	296.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,089	3.5	221.4
7	Alzheimer disease (G30)	966	3.1	196.4
8	Septicemia (A40-A41)	665	2.2	135.2
9	Influenza and pneumonia (J09-J18)	641	2.1	130.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	581	1.9	118.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	561	1.8	114.0
12	Parkinson disease (G20-G21)	387	1.3	78.7
13	Pneumonitis due to solids and liquids (J69)	261	0.8	53.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	169	0.5	34.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	125	0.4	25.4
...	All other causes (residual)	4,966	16.2	1,009.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 85 years and over				
...	All causes	19,462	100.0	13,164.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,352	27.5	3,620.3
2	Malignant neoplasms (C00-C97)	3,229	16.6	2,184.2
3	Cerebrovascular diseases (I60-I69)	1,350	6.9	913.2
4	Alzheimer disease (G30)	1,150	5.9	777.9
5	Chronic lower respiratory diseases (J40-J47)	893	4.6	604.1
6	Diabetes mellitus (E10-E14)	681	3.5	460.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	672	3.5	454.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	471	2.4	318.6
8	Influenza and pneumonia (J09-J18)	471	2.4	318.6
10	Septicemia (A40-A41)	380	2.0	257.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	1.6	217.1
12	Parkinson disease (G20-G21)	258	1.3	174.5
13	Pneumonitis due to solids and liquids (J69)	245	1.3	165.7
14	Nutritional deficiencies (E40-E64)	111	0.6	75.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	100	0.5	67.6
...	All other causes (residual)	3,778	19.4	2,555.6
Non-Hispanic, black, female, all ages				
...	All causes	161,264	100.0	726.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,750	22.8	165.5
2	Malignant neoplasms (C00-C97)	34,611	21.5	155.8
3	Cerebrovascular diseases (I60-I69)	10,522	6.5	47.4
4	Diabetes mellitus (E10-E14)	7,304	4.5	32.9
5	Alzheimer disease (G30)	6,515	4.0	29.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,025	3.7	27.1
7	Chronic lower respiratory diseases (J40-J47)	5,612	3.5	25.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,987	3.1	22.5
9	Septicemia (A40-A41)	3,516	2.2	15.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,399	2.1	15.3
11	Influenza and pneumonia (J09-J18)	2,874	1.8	12.9
12	Certain conditions originating in the perinatal period (P00-P96)	1,557	1.0	7.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,265	0.8	5.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,226	0.8	5.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1,002	0.6	4.5
...	All other causes (residual)	34,099	21.1	153.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 1-4 years				
...	All causes	450	100.0	37.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	28.9	10.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	12.4	4.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	11.3	4.2
4	Malignant neoplasms (C00-C97)	19	4.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.6	*
6	Influenza and pneumonia (J09-J18)	12	2.7	*
6	Certain conditions originating in the perinatal period (P00-P96)	12	2.7	*
8	Cerebrovascular diseases (I60-I69)	10	2.2	*
9	Chronic lower respiratory diseases (J40-J47)	8	1.8	*
10	Septicemia (A40-A41)	5	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.9	*
13	Anemias (D50-D64)	3	0.7	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Meningitis (G00,G03)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
...	All other causes (residual)	113	25.1	9.4
Non-Hispanic, black, female, 5-14 years				
...	All causes	512	100.0	16.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	25.4	4.3
2	Malignant neoplasms (C00-C97)	63	12.3	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	10.7	1.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	6.8	1.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.7	0.8
6	Chronic lower respiratory diseases (J40-J47)	23	4.5	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	3.9	0.7
8	Influenza and pneumonia (J09-J18)	12	2.3	*
9	Cerebrovascular diseases (I60-I69)	10	2.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.8	*
11	Diabetes mellitus (E10-E14)	7	1.4	*
12	Septicemia (A40-A41)	6	1.2	*
13	Anemias (D50-D64)	5	1.0	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.8	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
...	All other causes (residual)	107	20.9	3.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,653	100.0	50.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	502	30.4	15.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	294	17.8	9.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	145	8.8	4.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	95	5.7	2.9
5	Malignant neoplasms (C00-C97)	92	5.6	2.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	58	3.5	1.8
7	Diabetes mellitus (E10-E14)	36	2.2	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	1.9	1.0
9	Chronic lower respiratory diseases (J40-J47)	30	1.8	0.9
10	Anemias (D50-D64)	27	1.6	0.8
11	Cerebrovascular diseases (I60-I69)	19	1.1	*
12	Influenza and pneumonia (J09-J18)	16	1.0	*
13	Septicemia (A40-A41)	13	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.5	*
...	All other causes (residual)	277	16.8	8.5
Non-Hispanic, black, female, 25-34 years				
...	All causes	3,532	100.0	107.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	799	22.6	24.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	385	10.9	11.7
3	Malignant neoplasms (C00-C97)	359	10.2	10.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	331	9.4	10.1
5	Pregnancy, childbirth and the puerperium (O00-O99)	161	4.6	4.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	141	4.0	4.3
7	Diabetes mellitus (E10-E14)	140	4.0	4.3
8	Human immunodeficiency virus (HIV) disease (B20-B24)	104	2.9	3.2
9	Cerebrovascular diseases (I60-I69)	75	2.1	2.3
10	Anemias (D50-D64)	56	1.6	1.7
11	Influenza and pneumonia (J09-J18)	50	1.4	1.5
12	Septicemia (A40-A41)	49	1.4	1.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	1.4	1.5
14	Chronic lower respiratory diseases (J40-J47)	48	1.4	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	1.0	1.0
...	All other causes (residual)	751	21.3	22.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 35-44 years				
...	All causes	6,005	100.0	212.4
1	Malignant neoplasms (C00-C97)	1,175	19.6	41.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,052	17.5	37.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	763	12.7	27.0
4	Diabetes mellitus (E10-E14)	266	4.4	9.4
5	Cerebrovascular diseases (I60-I69)	222	3.7	7.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	203	3.4	7.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	182	3.0	6.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	127	2.1	4.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	124	2.1	4.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	105	1.7	3.7
11	Pregnancy, childbirth and the puerperium (O00-O99)	104	1.7	3.7
12	Septicemia (A40-A41)	102	1.7	3.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	80	1.3	2.8
13	Chronic lower respiratory diseases (J40-J47)	80	1.3	2.8
15	Influenza and pneumonia (J09-J18)	74	1.2	2.6
...	All other causes (residual)	1,346	22.4	47.6
Non-Hispanic, black, female, 45-54 years				
...	All causes	13,045	100.0	458.4
1	Malignant neoplasms (C00-C97)	3,493	26.8	122.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,692	20.6	94.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,058	8.1	37.2
4	Diabetes mellitus (E10-E14)	615	4.7	21.6
5	Cerebrovascular diseases (I60-I69)	601	4.6	21.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	323	2.5	11.4
7	Chronic lower respiratory diseases (J40-J47)	317	2.4	11.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	298	2.3	10.5
9	Septicemia (A40-A41)	287	2.2	10.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	281	2.2	9.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	221	1.7	7.8
12	Influenza and pneumonia (J09-J18)	186	1.4	6.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	118	0.9	4.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	81	0.6	2.8
15	Anemias (D50-D64)	67	0.5	2.4
...	All other causes (residual)	2,407	18.5	84.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 55-64 years				
...	All causes	26,946	100.0	1,012.7
1	Malignant neoplasms (C00-C97)	8,429	31.3	316.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,935	22.0	223.0
3	Diabetes mellitus (E10-E14)	1,360	5.0	51.1
4	Cerebrovascular diseases (I60-I69)	1,355	5.0	50.9
5	Chronic lower respiratory diseases (J40-J47)	1,054	3.9	39.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	978	3.6	36.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	851	3.2	32.0
8	Septicemia (A40-A41)	623	2.3	23.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	486	1.8	18.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	455	1.7	17.1
11	Influenza and pneumonia (J09-J18)	405	1.5	15.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	294	1.1	11.0
13	Viral hepatitis (B15-B19)	109	0.4	4.1
14	Pneumonitis due to solids and liquids (J69)	106	0.4	4.0
15	Anemias (D50-D64)	94	0.3	3.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	0.3	3.5
...	All other causes (residual)	4,318	16.0	162.3
Non-Hispanic, black, female, 65-74 years				
...	All causes	31,625	100.0	1,920.5
1	Malignant neoplasms (C00-C97)	9,386	29.7	570.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,267	23.0	441.3
3	Cerebrovascular diseases (I60-I69)	1,974	6.2	119.9
4	Diabetes mellitus (E10-E14)	1,728	5.5	104.9
5	Chronic lower respiratory diseases (J40-J47)	1,402	4.4	85.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,149	3.6	69.8
7	Septicemia (A40-A41)	838	2.6	50.9
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	632	2.0	38.4
9	Influenza and pneumonia (J09-J18)	524	1.7	31.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	512	1.6	31.1
11	Alzheimer disease (G30)	434	1.4	26.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	224	0.7	13.6
13	Pneumonitis due to solids and liquids (J69)	171	0.5	10.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	133	0.4	8.1
15	Parkinson disease (G20-G21)	120	0.4	7.3
...	All other causes (residual)	5,131	16.2	311.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 75-84 years				
...	All causes	34,337	100.0	4,369.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,307	24.2	1,057.2
2	Malignant neoplasms (C00-C97)	7,320	21.3	931.6
3	Cerebrovascular diseases (I60-I69)	2,686	7.8	341.8
4	Alzheimer disease (G30)	1,802	5.2	229.3
5	Diabetes mellitus (E10-E14)	1,725	5.0	219.5
6	Chronic lower respiratory diseases (J40-J47)	1,438	4.2	183.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,291	3.8	164.3
8	Septicemia (A40-A41)	820	2.4	104.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	780	2.3	99.3
10	Influenza and pneumonia (J09-J18)	678	2.0	86.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	434	1.3	55.2
12	Parkinson disease (G20-G21)	258	0.8	32.8
13	Pneumonitis due to solids and liquids (J69)	220	0.6	28.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	177	0.5	22.5
15	Nutritional deficiencies (E40-E64)	116	0.3	14.8
...	All other causes (residual)	6,285	18.3	799.8
Non-Hispanic, black, female, 85 years and over				
...	All causes	40,183	100.0	11,646.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,929	27.2	3,167.6
2	Malignant neoplasms (C00-C97)	4,271	10.6	1,237.9
3	Alzheimer disease (G30)	4,208	10.5	1,219.6
4	Cerebrovascular diseases (I60-I69)	3,557	8.9	1,030.9
5	Diabetes mellitus (E10-E14)	1,424	3.5	412.7
6	Chronic lower respiratory diseases (J40-J47)	1,212	3.0	351.3
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,186	3.0	343.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,179	2.9	341.7
9	Influenza and pneumonia (J09-J18)	895	2.2	259.4
10	Septicemia (A40-A41)	749	1.9	217.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	510	1.3	147.8
12	Pneumonitis due to solids and liquids (J69)	336	0.8	97.4
13	Nutritional deficiencies (E40-E64)	272	0.7	78.8
14	Parkinson disease (G20-G21)	221	0.5	64.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	164	0.4	47.5
...	All other causes (residual)	9,070	22.6	2,628.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, all ages				
...	All causes	19,198	100.0	703.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,472	18.1	127.2
2	Malignant neoplasms (C00-C97)	3,257	17.0	119.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,229	11.6	81.7
4	Diabetes mellitus (E10-E14)	1,114	5.8	40.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,047	5.5	38.4
6	Chronic lower respiratory diseases (J40-J47)	939	4.9	34.4
7	Cerebrovascular diseases (I60-I69)	737	3.8	27.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	602	3.1	22.1
9	Influenza and pneumonia (J09-J18)	391	2.0	14.3
10	Alzheimer disease (G30)	376	2.0	13.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	333	1.7	12.2
12	Septicemia (A40-A41)	295	1.5	10.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	266	1.4	9.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	190	1.0	7.0
15	Certain conditions originating in the perinatal period (P00-P96)	117	0.6	4.3
...	All other causes (residual)	3,833	20.0	140.4
Non-Hispanic, American Indian or Alaska Native, both sexes, 1-4 years				
...	All causes	72	100.0	45.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	34.7	15.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	11.1	*
3	Influenza and pneumonia (J09-J18)	5	6.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	6.9	*
5	Malignant neoplasms (C00-C97)	4	5.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.8	*
6	Cerebrovascular diseases (I60-I69)	2	2.8	*
8	Acute bronchitis and bronchiolitis (J20-J21)	1	1.4	*
8	Hernia (K40-K46)	1	1.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.4	*
...	All other causes (residual)	18	25.0	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 5-14 years				
...	All causes	102	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	39.2	9.8
2	Malignant neoplasms (C00-C97)	16	15.7	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	11.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	5.9	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	3.9	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	2.9	*
7	Certain conditions originating in the perinatal period (P00-P96)	2	2.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
8	Diabetes mellitus (E10-E14)	1	1.0	*
8	Influenza and pneumonia (J09-J18)	1	1.0	*
8	Legal intervention (Y35,Y89.0)	1	1.0	*
...	All other causes (residual)	15	14.7	*
Non-Hispanic, American Indian or Alaska Native, both sexes, 15-24 years				
...	All causes	562	100.0	132.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	256	45.6	60.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	159	28.3	37.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	8.0	10.6
4	Malignant neoplasms (C00-C97)	16	2.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	1.8	*
6	Legal intervention (Y35,Y89.0)	5	0.9	*
7	Cerebrovascular diseases (I60-I69)	4	0.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.7	*
9	Diabetes mellitus (E10-E14)	3	0.5	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.5	*
11	Septicemia (A40-A41)	2	0.4	*
11	Influenza and pneumonia (J09-J18)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
...	All other causes (residual)	48	8.5	11.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 25-34 years				
...	All causes	1,194	100.0	290.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	459	38.4	111.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	184	15.4	44.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	8.0	23.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	85	7.1	20.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	5.0	14.6
6	Malignant neoplasms (C00-C97)	28	2.3	6.8
7	Diabetes mellitus (E10-E14)	14	1.2	*
8	Septicemia (A40-A41)	12	1.0	*
8	Influenza and pneumonia (J09-J18)	12	1.0	*
10	Cerebrovascular diseases (I60-I69)	10	0.8	*
11	Legal intervention (Y35,Y89.0)	9	0.8	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	8	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.4	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.3	*
15	Chronic lower respiratory diseases (J40-J47)	3	0.3	*
...	All other causes (residual)	205	17.2	49.9
Non-Hispanic, American Indian or Alaska Native, both sexes, 35-44 years				
...	All causes	1,409	100.0	428.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	421	29.9	127.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	189	13.4	57.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	135	9.6	41.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	7.6	32.5
5	Malignant neoplasms (C00-C97)	82	5.8	24.9
6	Diabetes mellitus (E10-E14)	75	5.3	22.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	3.8	16.4
8	Cerebrovascular diseases (I60-I69)	20	1.4	6.1
9	Septicemia (A40-A41)	16	1.1	*
9	Influenza and pneumonia (J09-J18)	16	1.1	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	10	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.5	*
15	Viral hepatitis (B15-B19)	4	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.3	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.3	*
...	All other causes (residual)	244	17.3	74.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 45-54 years				
...	All causes	2,286	100.0	686.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	389	17.0	116.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	359	15.7	107.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	311	13.6	93.4
4	Malignant neoplasms (C00-C97)	308	13.5	92.5
5	Diabetes mellitus (E10-E14)	136	5.9	40.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	79	3.5	23.7
7	Cerebrovascular diseases (I60-I69)	55	2.4	16.5
8	Chronic lower respiratory diseases (J40-J47)	45	2.0	13.5
9	Influenza and pneumonia (J09-J18)	43	1.9	12.9
10	Septicemia (A40-A41)	42	1.8	12.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.6	11.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	1.6	11.1
13	Viral hepatitis (B15-B19)	17	0.7	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.3	*
...	All other causes (residual)	403	17.6	121.1
Non-Hispanic, American Indian or Alaska Native, both sexes, 55-64 years				
...	All causes	3,543	100.0	1,099.3
1	Malignant neoplasms (C00-C97)	875	24.7	271.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	658	18.6	204.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	283	8.0	87.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	278	7.8	86.3
5	Diabetes mellitus (E10-E14)	244	6.9	75.7
6	Chronic lower respiratory diseases (J40-J47)	178	5.0	55.2
7	Cerebrovascular diseases (I60-I69)	102	2.9	31.6
8	Influenza and pneumonia (J09-J18)	64	1.8	19.9
9	Septicemia (A40-A41)	61	1.7	18.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	1.6	17.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	1.1	11.8
12	Viral hepatitis (B15-B19)	34	1.0	10.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	0.9	9.6
14	Pneumonitis due to solids and liquids (J69)	23	0.6	7.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	0.4	*
...	All other causes (residual)	603	17.0	187.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 65-74 years				
...	All causes	3,744	100.0	1,937.0
1	Malignant neoplasms (C00-C97)	958	25.6	495.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	766	20.5	396.3
3	Diabetes mellitus (E10-E14)	292	7.8	151.1
4	Chronic lower respiratory diseases (J40-J47)	283	7.6	146.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	4.0	77.1
6	Cerebrovascular diseases (I60-I69)	144	3.8	74.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	141	3.8	72.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	95	2.5	49.1
9	Influenza and pneumonia (J09-J18)	74	2.0	38.3
10	Septicemia (A40-A41)	70	1.9	36.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	1.1	22.2
12	Alzheimer disease (G30)	25	0.7	12.9
13	Parkinson disease (G20-G21)	18	0.5	*
14	Pneumonitis due to solids and liquids (J69)	17	0.5	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	0.4	*
...	All other causes (residual)	654	17.5	338.4
Non-Hispanic, American Indian or Alaska Native, both sexes, 75-84 years				
...	All causes	3,529	100.0	4,423.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	803	22.8	1,006.6
2	Malignant neoplasms (C00-C97)	689	19.5	863.7
3	Chronic lower respiratory diseases (J40-J47)	264	7.5	330.9
4	Diabetes mellitus (E10-E14)	246	7.0	308.4
5	Cerebrovascular diseases (I60-I69)	218	6.2	273.3
6	Alzheimer disease (G30)	152	4.3	190.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	3.2	141.6
8	Influenza and pneumonia (J09-J18)	96	2.7	120.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	2.3	101.5
10	Septicemia (A40-A41)	60	1.7	75.2
11	Parkinson disease (G20-G21)	59	1.7	74.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	53	1.5	66.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	36	1.0	45.1
14	Pneumonitis due to solids and liquids (J69)	26	0.7	32.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.4	*
...	All other causes (residual)	618	17.5	774.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 85 years and over				
...	All causes	2,473	100.0	9,157.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	666	26.9	2,466.1
2	Malignant neoplasms (C00-C97)	281	11.4	1,040.5
3	Alzheimer disease (G30)	193	7.8	714.7
4	Cerebrovascular diseases (I60-I69)	182	7.4	673.9
5	Chronic lower respiratory diseases (J40-J47)	155	6.3	573.9
6	Diabetes mellitus (E10-E14)	103	4.2	381.4
7	Influenza and pneumonia (J09-J18)	77	3.1	285.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	2.8	259.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.9	170.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	1.7	151.8
11	Parkinson disease (G20-G21)	36	1.5	133.3
12	Septicemia (A40-A41)	29	1.2	107.4
13	Nutritional deficiencies (E40-E64)	20	0.8	74.1
14	Pneumonitis due to solids and liquids (J69)	18	0.7	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	14	0.6	*
...	All other causes (residual)	542	21.9	2,007.0
Non-Hispanic, American Indian or Alaska Native, male, all ages				
...	All causes	10,502	100.0	784.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,033	19.4	151.8
2	Malignant neoplasms (C00-C97)	1,725	16.4	128.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,448	13.8	108.1
4	Diabetes mellitus (E10-E14)	624	5.9	46.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	559	5.3	41.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	448	4.3	33.5
7	Chronic lower respiratory diseases (J40-J47)	441	4.2	32.9
8	Cerebrovascular diseases (I60-I69)	322	3.1	24.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	199	1.9	14.9
10	Influenza and pneumonia (J09-J18)	186	1.8	13.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	1.5	11.5
12	Septicemia (A40-A41)	130	1.2	9.7
13	Alzheimer disease (G30)	120	1.1	9.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	0.9	7.2
15	Parkinson disease (G20-G21)	68	0.6	5.1
...	All other causes (residual)	1,948	18.5	145.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 1-4 years				
...	All causes	33	100.0	40.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	33.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	12.1	*
3	Malignant neoplasms (C00-C97)	3	9.1	*
4	Cerebrovascular diseases (I60-I69)	2	6.1	*
4	Influenza and pneumonia (J09-J18)	2	6.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	6.1	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	3.0	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	3.0	*
7	Hernia (K40-K46)	1	3.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	3.0	*
...	All other causes (residual)	5	15.2	*
Non-Hispanic, American Indian or Alaska Native, male, 5-14 years				
...	All causes	55	100.0	26.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	41.8	11.1
2	Malignant neoplasms (C00-C97)	9	16.4	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	9.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	7.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	3.6	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	1.8	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.8	*
6	Legal intervention (Y35,Y89.0)	1	1.8	*
...	All other causes (residual)	8	14.5	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 15-24 years				
...	All causes	392	100.0	181.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	45.2	82.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	29.6	53.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	9.7	17.6
4	Malignant neoplasms (C00-C97)	11	2.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.0	*
6	Legal intervention (Y35,Y89.0)	5	1.3	*
7	Diabetes mellitus (E10-E14)	2	0.5	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
9	Septicemia (A40-A41)	1	0.3	*
9	Influenza and pneumonia (J09-J18)	1	0.3	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (residual)	29	7.4	13.4
Non-Hispanic, American Indian or Alaska Native, male, 25-34 years				
...	All causes	779	100.0	376.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	308	39.5	148.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	130	16.7	62.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	8.7	32.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	46	5.9	22.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	5.1	19.3
6	Malignant neoplasms (C00-C97)	11	1.4	*
7	Diabetes mellitus (E10-E14)	10	1.3	*
8	Legal intervention (Y35,Y89.0)	9	1.2	*
9	Septicemia (A40-A41)	7	0.9	*
10	Influenza and pneumonia (J09-J18)	6	0.8	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.6	*
11	Cerebrovascular diseases (I60-I69)	5	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.3	*
15	Meningitis (G00,G03)	1	0.1	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.1	*
15	Aortic aneurysm and dissection (I71)	1	0.1	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.1	*
15	Pneumonitis due to solids and liquids (J69)	1	0.1	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.1	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.1	*
...	All other causes (residual)	122	15.7	59.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 35-44 years				
...	All causes	871	100.0	538.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	274	31.5	169.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	98	11.3	60.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	84	9.6	51.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	9.1	48.8
5	Diabetes mellitus (E10-E14)	46	5.3	28.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	4.8	26.0
7	Malignant neoplasms (C00-C97)	35	4.0	21.6
8	Cerebrovascular diseases (I60-I69)	14	1.6	*
9	Influenza and pneumonia (J09-J18)	12	1.4	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.1	*
11	Septicemia (A40-A41)	8	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.8	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.5	*
14	Viral hepatitis (B15-B19)	3	0.3	*
14	Nutritional deficiencies (E40-E64)	3	0.3	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.3	*
14	Aortic aneurysm and dissection (I71)	3	0.3	*
14	Chronic lower respiratory diseases (J40-J47)	3	0.3	*
14	Legal intervention (Y35,Y89.0)	3	0.3	*
...	All other causes (residual)	140	16.1	86.6
Non-Hispanic, American Indian or Alaska Native, male, 45-54 years				
...	All causes	1,367	100.0	848.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	251	18.4	155.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	249	18.2	154.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	167	12.2	103.7
4	Malignant neoplasms (C00-C97)	143	10.5	88.8
5	Diabetes mellitus (E10-E14)	83	6.1	51.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	4.5	38.5
7	Cerebrovascular diseases (I60-I69)	33	2.4	20.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	2.2	18.6
9	Septicemia (A40-A41)	23	1.7	14.3
10	Influenza and pneumonia (J09-J18)	22	1.6	13.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.4	*
12	Chronic lower respiratory diseases (J40-J47)	18	1.3	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	1.0	*
14	Viral hepatitis (B15-B19)	12	0.9	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.4	*
...	All other causes (residual)	235	17.2	145.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 55-64 years				
...	All causes	2,068	100.0	1,370.5
1	Malignant neoplasms (C00-C97)	491	23.7	325.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	423	20.5	280.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	199	9.6	131.9
4	Diabetes mellitus (E10-E14)	164	7.9	108.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	153	7.4	101.4
6	Chronic lower respiratory diseases (J40-J47)	81	3.9	53.7
7	Cerebrovascular diseases (I60-I69)	49	2.4	32.5
8	Influenza and pneumonia (J09-J18)	35	1.7	23.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	1.5	19.9
10	Septicemia (A40-A41)	29	1.4	19.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.3	17.9
12	Viral hepatitis (B15-B19)	22	1.1	14.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.9	*
14	Pneumonitis due to solids and liquids (J69)	10	0.5	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	0.5	*
...	All other causes (residual)	327	15.8	216.7
Non-Hispanic, American Indian or Alaska Native, male, 65-74 years				
...	All causes	2,084	100.0	2,322.1
1	Malignant neoplasms (C00-C97)	528	25.3	588.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	495	23.8	551.6
3	Diabetes mellitus (E10-E14)	162	7.8	180.5
4	Chronic lower respiratory diseases (J40-J47)	139	6.7	154.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	4.3	100.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	85	4.1	94.7
7	Cerebrovascular diseases (I60-I69)	76	3.6	84.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	2.4	55.7
9	Influenza and pneumonia (J09-J18)	37	1.8	41.2
10	Septicemia (A40-A41)	29	1.4	32.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	0.9	*
12	Pneumonitis due to solids and liquids (J69)	13	0.6	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	0.6	*
14	Parkinson disease (G20-G21)	12	0.6	*
15	Alzheimer disease (G30)	10	0.5	*
...	All other causes (residual)	326	15.6	363.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 75-84 years				
...	All causes	1,762	100.0	5,051.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	443	25.1	1,270.1
2	Malignant neoplasms (C00-C97)	373	21.2	1,069.4
3	Chronic lower respiratory diseases (J40-J47)	135	7.7	387.1
4	Diabetes mellitus (E10-E14)	121	6.9	346.9
5	Cerebrovascular diseases (I60-I69)	83	4.7	238.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	3.9	195.0
7	Alzheimer disease (G30)	56	3.2	160.6
8	Influenza and pneumonia (J09-J18)	42	2.4	120.4
9	Parkinson disease (G20-G21)	37	2.1	106.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	1.8	91.7
11	Septicemia (A40-A41)	27	1.5	77.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	1.5	77.4
13	Pneumonitis due to solids and liquids (J69)	14	0.8	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	9	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.5	*
...	All other causes (residual)	287	16.3	822.9
Non-Hispanic, American Indian or Alaska Native, male, 85 years and over				
...	All causes	940	100.0	9,649.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	286	30.4	2,936.0
2	Malignant neoplasms (C00-C97)	121	12.9	1,242.2
3	Chronic lower respiratory diseases (J40-J47)	63	6.7	646.8
4	Cerebrovascular diseases (I60-I69)	60	6.4	616.0
5	Alzheimer disease (G30)	51	5.4	523.6
6	Diabetes mellitus (E10-E14)	36	3.8	369.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	3.4	328.5
8	Influenza and pneumonia (J09-J18)	28	3.0	287.4
9	Parkinson disease (G20-G21)	19	2.0	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	1.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.5	*
12	Pneumonitis due to solids and liquids (J69)	10	1.1	*
13	Nutritional deficiencies (E40-E64)	8	0.9	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	8	0.9	*
15	Septicemia (A40-A41)	5	0.5	*
15	Pneumoconioses and chemical effects (J60-J66,J68)	5	0.5	*
...	All other causes (residual)	179	19.0	1,837.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, all ages				
...	All causes	8,696	100.0	625.5
1	Malignant neoplasms (C00-C97)	1,532	17.6	110.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,439	16.5	103.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	781	9.0	56.2
4	Chronic lower respiratory diseases (J40-J47)	498	5.7	35.8
5	Diabetes mellitus (E10-E14)	490	5.6	35.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	488	5.6	35.1
7	Cerebrovascular diseases (I60-I69)	415	4.8	29.9
8	Alzheimer disease (G30)	256	2.9	18.4
9	Influenza and pneumonia (J09-J18)	205	2.4	14.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	179	2.1	12.9
11	Septicemia (A40-A41)	165	1.9	11.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	154	1.8	11.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	93	1.1	6.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	0.8	4.8
15	Certain conditions originating in the perinatal period (P00-P96)	56	0.6	4.0
...	All other causes (residual)	1,878	21.6	135.1
Non-Hispanic, American Indian or Alaska Native, female, 1-4 years				
...	All causes	39	100.0	50.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	35.9	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	15.4	*
3	Influenza and pneumonia (J09-J18)	3	7.7	*
4	Malignant neoplasms (C00-C97)	1	2.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.6	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.6	*
...	All other causes (residual)	13	33.3	*
Non-Hispanic, American Indian or Alaska Native, female, 5-14 years				
...	All causes	47	100.0	23.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	36.2	*
2	Malignant neoplasms (C00-C97)	7	14.9	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	14.9	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	6.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.3	*
6	Diabetes mellitus (E10-E14)	1	2.1	*
6	Influenza and pneumonia (J09-J18)	1	2.1	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	2.1	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	2.1	*
...	All other causes (residual)	7	14.9	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 15-24 years				
...	All causes	170	100.0	81.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	46.5	37.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	25.3	20.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	4.1	*
4	Malignant neoplasms (C00-C97)	5	2.9	*
5	Cerebrovascular diseases (I60-I69)	4	2.4	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.8	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.2	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.2	*
9	Septicemia (A40-A41)	1	0.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.6	*
9	Diabetes mellitus (E10-E14)	1	0.6	*
9	Influenza and pneumonia (J09-J18)	1	0.6	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
...	All other causes (residual)	19	11.2	*
Non-Hispanic, American Indian or Alaska Native, female, 25-34 years				
...	All causes	415	100.0	203.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	151	36.4	73.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	13.0	26.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	39	9.4	19.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	6.7	13.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	4.8	9.8
6	Malignant neoplasms (C00-C97)	17	4.1	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	8	1.9	*
8	Influenza and pneumonia (J09-J18)	6	1.4	*
9	Septicemia (A40-A41)	5	1.2	*
9	Cerebrovascular diseases (I60-I69)	5	1.2	*
11	Diabetes mellitus (E10-E14)	4	1.0	*
12	Nutritional deficiencies (E40-E64)	2	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (residual)	69	16.6	33.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 35-44 years				
...	All causes	538	100.0	321.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	27.3	87.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	91	16.9	54.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	10.4	33.5
4	Malignant neoplasms (C00-C97)	47	8.7	28.1
5	Diabetes mellitus (E10-E14)	29	5.4	17.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	4.3	13.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	2.2	*
8	Septicemia (A40-A41)	8	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.3	*
10	Cerebrovascular diseases (I60-I69)	6	1.1	*
11	Influenza and pneumonia (J09-J18)	4	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.6	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	2	0.4	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.4	*
...	All other causes (residual)	91	16.9	54.4
Non-Hispanic, American Indian or Alaska Native, female, 45-54 years				
...	All causes	919	100.0	535.0
1	Malignant neoplasms (C00-C97)	165	18.0	96.0
2	Chronic liver disease and cirrhosis (K70,K73-K74)	144	15.7	83.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	15.0	80.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	110	12.0	64.0
5	Diabetes mellitus (E10-E14)	53	5.8	30.9
6	Chronic lower respiratory diseases (J40-J47)	27	2.9	15.7
7	Cerebrovascular diseases (I60-I69)	22	2.4	12.8
8	Influenza and pneumonia (J09-J18)	21	2.3	12.2
9	Septicemia (A40-A41)	19	2.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	2.0	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	1.8	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	0.8	*
13	Viral hepatitis (B15-B19)	5	0.5	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.3	*
14	Pneumonitis due to solids and liquids (J69)	3	0.3	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (residual)	161	17.5	93.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 55-64 years				
...	All causes	1,475	100.0	860.6
1	Malignant neoplasms (C00-C97)	384	26.0	224.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	235	15.9	137.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	125	8.5	72.9
4	Chronic lower respiratory diseases (J40-J47)	97	6.6	56.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	5.7	49.0
6	Diabetes mellitus (E10-E14)	80	5.4	46.7
7	Cerebrovascular diseases (I60-I69)	53	3.6	30.9
8	Septicemia (A40-A41)	32	2.2	18.7
9	Influenza and pneumonia (J09-J18)	29	2.0	16.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	2.0	16.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.9	*
11	Pneumonitis due to solids and liquids (J69)	13	0.9	*
13	Viral hepatitis (B15-B19)	12	0.8	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.6	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	0.5	*
...	All other causes (residual)	272	18.4	158.7
Non-Hispanic, American Indian or Alaska Native, female, 65-74 years				
...	All causes	1,660	100.0	1,603.2
1	Malignant neoplasms (C00-C97)	430	25.9	415.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	271	16.3	261.7
3	Chronic lower respiratory diseases (J40-J47)	144	8.7	139.1
4	Diabetes mellitus (E10-E14)	130	7.8	125.6
5	Cerebrovascular diseases (I60-I69)	68	4.1	65.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	3.6	57.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	56	3.4	54.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	2.7	43.5
9	Septicemia (A40-A41)	41	2.5	39.6
10	Influenza and pneumonia (J09-J18)	37	2.2	35.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	1.4	23.2
12	Alzheimer disease (G30)	15	0.9	*
13	Parkinson disease (G20-G21)	6	0.4	*
14	Viral hepatitis (B15-B19)	5	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.3	*
14	Nutritional deficiencies (E40-E64)	5	0.3	*
14	Aortic aneurysm and dissection (I71)	5	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (residual)	309	18.6	298.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 75-84 years				
...	All causes	1,767	100.0	3,935.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	360	20.4	801.8
2	Malignant neoplasms (C00-C97)	316	17.9	703.8
3	Cerebrovascular diseases (I60-I69)	135	7.6	300.7
4	Chronic lower respiratory diseases (J40-J47)	129	7.3	287.3
5	Diabetes mellitus (E10-E14)	125	7.1	278.4
6	Alzheimer disease (G30)	96	5.4	213.8
7	Influenza and pneumonia (J09-J18)	54	3.1	120.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	2.8	109.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	2.5	100.2
10	Septicemia (A40-A41)	33	1.9	73.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	27	1.5	60.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	1.5	57.9
13	Parkinson disease (G20-G21)	22	1.2	49.0
14	Pneumonitis due to solids and liquids (J69)	12	0.7	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	9	0.5	*
...	All other causes (residual)	329	18.6	732.8
Non-Hispanic, American Indian or Alaska Native, female, 85 years and over				
...	All causes	1,533	100.0	8,879.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	380	24.8	2,201.0
2	Malignant neoplasms (C00-C97)	160	10.4	926.7
3	Alzheimer disease (G30)	142	9.3	822.5
4	Cerebrovascular diseases (I60-I69)	122	8.0	706.6
5	Chronic lower respiratory diseases (J40-J47)	92	6.0	532.9
6	Diabetes mellitus (E10-E14)	67	4.4	388.1
7	Influenza and pneumonia (J09-J18)	49	3.2	283.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	2.5	220.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	2.1	185.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	1.7	150.6
11	Septicemia (A40-A41)	24	1.6	139.0
12	Parkinson disease (G20-G21)	17	1.1	*
13	Nutritional deficiencies (E40-E64)	12	0.8	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	11	0.7	*
15	Pneumonitis due to solids and liquids (J69)	8	0.5	*
...	All other causes (residual)	353	23.0	2,044.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, all ages				
...	All causes	72,598	100.0	359.8
1	Malignant neoplasms (C00-C97)	18,257	25.1	90.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,490	21.3	76.8
3	Cerebrovascular diseases (I60-I69)	5,430	7.5	26.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,256	4.5	16.1
5	Diabetes mellitus (E10-E14)	3,047	4.2	15.1
6	Alzheimer disease (G30)	2,583	3.6	12.8
7	Influenza and pneumonia (J09-J18)	2,286	3.1	11.3
8	Chronic lower respiratory diseases (J40-J47)	2,063	2.8	10.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,534	2.1	7.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,472	2.0	7.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,414	1.9	7.0
12	Septicemia (A40-A41)	879	1.2	4.4
13	Parkinson disease (G20-G21)	869	1.2	4.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	725	1.0	3.6
15	Pneumonitis due to solids and liquids (J69)	549	0.8	2.7
...	All other causes (residual)	12,744	17.6	63.2
Non-Hispanic, Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	148	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	24.3	3.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	14.2	2.2
3	Malignant neoplasms (C00-C97)	20	13.5	2.1
4	Influenza and pneumonia (J09-J18)	6	4.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	2.7	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	2.7	*
7	Septicemia (A40-A41)	3	2.0	*
7	Anemias (D50-D64)	3	2.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	3	2.0	*
10	Meningitis (G00,G03)	2	1.4	*
10	Cerebrovascular diseases (I60-I69)	2	1.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.7	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.7	*
12	Hernia (K40-K46)	1	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.7	*
...	All other causes (residual)	40	27.0	4.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	208	100.0	8.7
1	Malignant neoplasms (C00-C97)	42	20.2	1.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	17.8	1.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	10.6	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	10.1	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	4.3	*
6	Septicemia (A40-A41)	6	2.9	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	1.9	*
8	Anemias (D50-D64)	3	1.4	*
8	Influenza and pneumonia (J09-J18)	3	1.4	*
8	Chronic lower respiratory diseases (J40-J47)	3	1.4	*
11	Cerebrovascular diseases (I60-I69)	2	1.0	*
11	Acute bronchitis and bronchiolitis (J20-J21)	2	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	1.0	*
14	Meningococcal infection (A39)	1	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
14	Diabetes mellitus (E10-E14)	1	0.5	*
14	Meningitis (G00,G03)	1	0.5	*
...	All other causes (residual)	48	23.1	2.0
Non-Hispanic, Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	960	100.0	36.0
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	315	32.8	11.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	283	29.5	10.6
3	Malignant neoplasms (C00-C97)	79	8.2	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	6.1	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.9	1.4
6	Influenza and pneumonia (J09-J18)	15	1.6	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	1.3	*
8	Cerebrovascular diseases (I60-I69)	6	0.6	*
9	Septicemia (A40-A41)	4	0.4	*
9	Diabetes mellitus (E10-E14)	4	0.4	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.4	*
12	Aortic aneurysm and dissection (I71)	3	0.3	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.3	*
12	Pneumonitis due to solids and liquids (J69)	3	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (residual)	130	13.5	4.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	1,500	100.0	43.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	433	28.9	12.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	299	19.9	8.6
3	Malignant neoplasms (C00-C97)	206	13.7	5.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	6.5	2.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	5.4	2.3
6	Cerebrovascular diseases (I60-I69)	34	2.3	1.0
7	Diabetes mellitus (E10-E14)	19	1.3	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.3	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	18	1.2	*
10	Influenza and pneumonia (J09-J18)	16	1.1	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.7	*
12	Septicemia (A40-A41)	9	0.6	*
12	Aortic aneurysm and dissection (I71)	9	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
...	All other causes (residual)	226	15.1	6.5
Non-Hispanic, Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	2,227	100.0	70.2
1	Malignant neoplasms (C00-C97)	605	27.2	19.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	323	14.5	10.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	310	13.9	9.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	212	9.5	6.7
5	Cerebrovascular diseases (I60-I69)	128	5.7	4.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	68	3.1	2.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	2.7	1.9
8	Diabetes mellitus (E10-E14)	57	2.6	1.8
9	Influenza and pneumonia (J09-J18)	25	1.1	0.8
10	Septicemia (A40-A41)	21	0.9	0.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21	0.9	0.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.9	0.7
13	Viral hepatitis (B15-B19)	14	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	14	0.6	*
...	All other causes (residual)	333	15.0	10.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	4,685	100.0	171.7
1	Malignant neoplasms (C00-C97)	1,785	38.1	65.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	800	17.1	29.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	353	7.5	12.9
4	Cerebrovascular diseases (I60-I69)	275	5.9	10.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	207	4.4	7.6
6	Diabetes mellitus (E10-E14)	182	3.9	6.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	130	2.8	4.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	1.2	2.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	51	1.1	1.9
10	Influenza and pneumonia (J09-J18)	50	1.1	1.8
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	1.0	1.8
12	Septicemia (A40-A41)	40	0.9	1.5
13	Chronic lower respiratory diseases (J40-J47)	38	0.8	1.4
14	Aortic aneurysm and dissection (I71)	32	0.7	1.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	29	0.6	1.1
...	All other causes (residual)	608	13.0	22.3
Non-Hispanic, Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	8,502	100.0	389.5
1	Malignant neoplasms (C00-C97)	3,474	40.9	159.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,670	19.6	76.5
3	Cerebrovascular diseases (I60-I69)	501	5.9	23.0
4	Diabetes mellitus (E10-E14)	442	5.2	20.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	351	4.1	16.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	2.0	7.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	157	1.8	7.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	154	1.8	7.1
9	Influenza and pneumonia (J09-J18)	130	1.5	6.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	122	1.4	5.6
11	Septicemia (A40-A41)	115	1.4	5.3
11	Chronic lower respiratory diseases (J40-J47)	115	1.4	5.3
13	Viral hepatitis (B15-B19)	65	0.8	3.0
14	Aortic aneurysm and dissection (I71)	51	0.6	2.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.5	2.0
...	All other causes (residual)	945	11.1	43.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	12,893	100.0	894.9
1	Malignant neoplasms (C00-C97)	4,608	35.7	319.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,710	21.0	188.1
3	Cerebrovascular diseases (I60-I69)	874	6.8	60.7
4	Diabetes mellitus (E10-E14)	674	5.2	46.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	382	3.0	26.5
6	Chronic lower respiratory diseases (J40-J47)	341	2.6	23.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	292	2.3	20.3
8	Influenza and pneumonia (J09-J18)	290	2.2	20.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	196	1.5	13.6
10	Septicemia (A40-A41)	178	1.4	12.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	161	1.2	11.2
12	Parkinson disease (G20-G21)	140	1.1	9.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	0.9	8.1
14	Alzheimer disease (G30)	93	0.7	6.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	88	0.7	6.1
...	All other causes (residual)	1,750	13.6	121.5
Non-Hispanic, Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	17,530	100.0	2,583.9
1	Malignant neoplasms (C00-C97)	4,454	25.4	656.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,890	22.2	573.4
3	Cerebrovascular diseases (I60-I69)	1,450	8.3	213.7
4	Diabetes mellitus (E10-E14)	886	5.1	130.6
5	Chronic lower respiratory diseases (J40-J47)	688	3.9	101.4
6	Influenza and pneumonia (J09-J18)	579	3.3	85.3
7	Alzheimer disease (G30)	558	3.2	82.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	502	2.9	74.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	464	2.6	68.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	373	2.1	55.0
11	Parkinson disease (G20-G21)	348	2.0	51.3
12	Septicemia (A40-A41)	244	1.4	36.0
13	Pneumonitis due to solids and liquids (J69)	153	0.9	22.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	128	0.7	18.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	115	0.7	17.0
...	All other causes (residual)	2,698	15.4	397.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	22,988	100.0	8,567.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,928	25.8	2,209.4
2	Malignant neoplasms (C00-C97)	2,980	13.0	1,110.7
3	Cerebrovascular diseases (I60-I69)	2,154	9.4	802.8
4	Alzheimer disease (G30)	1,916	8.3	714.1
5	Influenza and pneumonia (J09-J18)	1,164	5.1	433.8
6	Chronic lower respiratory diseases (J40-J47)	855	3.7	318.7
7	Diabetes mellitus (E10-E14)	782	3.4	291.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	702	3.1	261.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	536	2.3	199.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	519	2.3	193.4
11	Parkinson disease (G20-G21)	356	1.5	132.7
12	Pneumonitis due to solids and liquids (J69)	292	1.3	108.8
13	Septicemia (A40-A41)	252	1.1	93.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	127	0.6	47.3
15	Nutritional deficiencies (E40-E64)	113	0.5	42.1
...	All other causes (residual)	4,312	18.8	1,607.1
Non-Hispanic, Asian or Pacific Islander, male, all ages				
...	All causes	37,236	100.0	386.2
1	Malignant neoplasms (C00-C97)	9,243	24.8	95.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,418	22.6	87.3
3	Cerebrovascular diseases (I60-I69)	2,442	6.6	25.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,074	5.6	21.5
5	Diabetes mellitus (E10-E14)	1,612	4.3	16.7
6	Chronic lower respiratory diseases (J40-J47)	1,198	3.2	12.4
7	Influenza and pneumonia (J09-J18)	1,155	3.1	12.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	990	2.7	10.3
9	Alzheimer disease (G30)	790	2.1	8.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	789	2.1	8.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	654	1.8	6.8
12	Parkinson disease (G20-G21)	489	1.3	5.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	468	1.3	4.9
14	Septicemia (A40-A41)	460	1.2	4.8
15	Pneumonitis due to solids and liquids (J69)	317	0.9	3.3
...	All other causes (residual)	6,137	16.5	63.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 1-4 years				
...	All causes	96	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	27.1	5.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	14.6	*
3	Malignant neoplasms (C00-C97)	13	13.5	*
4	Influenza and pneumonia (J09-J18)	3	3.1	*
5	Septicemia (A40-A41)	2	2.1	*
5	Meningitis (G00,G03)	2	2.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.1	*
5	Cerebrovascular diseases (I60-I69)	2	2.1	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.1	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
...	All other causes (residual)	28	29.2	5.7
Non-Hispanic, Asian or Pacific Islander, male, 5-14 years				
...	All causes	114	100.0	9.4
1	Malignant neoplasms (C00-C97)	23	20.2	1.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	15.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	10.5	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	9.6	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	6.1	*
6	Influenza and pneumonia (J09-J18)	3	2.6	*
7	Septicemia (A40-A41)	2	1.8	*
7	Anemias (D50-D64)	2	1.8	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.8	*
7	Certain conditions originating in the perinatal period (P00-P96)	2	1.8	*
11	Meningococcal infection (A39)	1	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
11	Meningitis (G00,G03)	1	0.9	*
11	Acute bronchitis and bronchiolitis (J20-J21)	1	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.9	*
...	All other causes (residual)	27	23.7	2.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 15-24 years				
...	All causes	667	100.0	49.6
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	228	34.2	16.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	30.4	15.1
3	Malignant neoplasms (C00-C97)	48	7.2	3.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	7.2	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.3	1.6
6	Influenza and pneumonia (J09-J18)	11	1.6	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.3	*
8	Cerebrovascular diseases (I60-I69)	6	0.9	*
9	Septicemia (A40-A41)	3	0.4	*
9	Aortic aneurysm and dissection (I71)	3	0.4	*
11	Anemias (D50-D64)	2	0.3	*
11	Diabetes mellitus (E10-E14)	2	0.3	*
11	Pneumonitis due to solids and liquids (J69)	2	0.3	*
11	Legal intervention (Y35,Y89.0)	2	0.3	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	76	11.4	5.6
Non-Hispanic, Asian or Pacific Islander, male, 25-34 years				
...	All causes	1,013	100.0	60.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	349	34.5	20.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	225	22.2	13.3
3	Malignant neoplasms (C00-C97)	80	7.9	4.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	7.3	4.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	5.5	3.3
6	Cerebrovascular diseases (I60-I69)	20	2.0	1.2
7	Diabetes mellitus (E10-E14)	14	1.4	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	12	1.2	*
9	Aortic aneurysm and dissection (I71)	9	0.9	*
9	Influenza and pneumonia (J09-J18)	9	0.9	*
11	Septicemia (A40-A41)	6	0.6	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.5	*
13	Legal intervention (Y35,Y89.0)	5	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (residual)	135	13.3	8.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 35-44 years				
...	All causes	1,374	100.0	92.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	259	18.9	17.4
2	Malignant neoplasms (C00-C97)	253	18.4	17.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	17.6	16.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	141	10.3	9.5
5	Cerebrovascular diseases (I60-I69)	75	5.5	5.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	49	3.6	3.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	3.1	2.8
8	Diabetes mellitus (E10-E14)	32	2.3	2.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	1.4	*
10	Influenza and pneumonia (J09-J18)	15	1.1	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.9	*
12	Viral hepatitis (B15-B19)	12	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.8	*
14	Aortic aneurysm and dissection (I71)	10	0.7	*
15	Septicemia (A40-A41)	6	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.4	*
...	All other causes (residual)	189	13.8	12.7
Non-Hispanic, Asian or Pacific Islander, male, 45-54 years				
...	All causes	2,794	100.0	219.3
1	Malignant neoplasms (C00-C97)	796	28.5	62.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	620	22.2	48.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	252	9.0	19.8
4	Cerebrovascular diseases (I60-I69)	154	5.5	12.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	138	4.9	10.8
6	Diabetes mellitus (E10-E14)	114	4.1	8.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	106	3.8	8.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.5	3.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	1.3	2.9
10	Influenza and pneumonia (J09-J18)	33	1.2	2.6
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	1.2	2.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	29	1.0	2.3
13	Aortic aneurysm and dissection (I71)	26	0.9	2.0
14	Viral hepatitis (B15-B19)	23	0.8	1.8
15	Septicemia (A40-A41)	22	0.8	1.7
15	Chronic lower respiratory diseases (J40-J47)	22	0.8	1.7
...	All other causes (residual)	348	12.5	27.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 55-64 years				
...	All causes	5,031	100.0	506.3
1	Malignant neoplasms (C00-C97)	1,734	34.5	174.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,207	24.0	121.5
3	Cerebrovascular diseases (I60-I69)	299	5.9	30.1
4	Diabetes mellitus (E10-E14)	287	5.7	28.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	237	4.7	23.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	2.3	11.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	112	2.2	11.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	2.1	10.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	80	1.6	8.1
10	Influenza and pneumonia (J09-J18)	74	1.5	7.4
11	Septicemia (A40-A41)	69	1.4	6.9
12	Chronic lower respiratory diseases (J40-J47)	64	1.3	6.4
13	Viral hepatitis (B15-B19)	43	0.9	4.3
14	Aortic aneurysm and dissection (I71)	32	0.6	3.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	0.6	3.0
...	All other causes (residual)	543	10.8	54.6
Non-Hispanic, Asian or Pacific Islander, male, 65-74 years				
...	All causes	7,352	100.0	1,150.4
1	Malignant neoplasms (C00-C97)	2,475	33.7	387.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,731	23.5	270.9
3	Cerebrovascular diseases (I60-I69)	455	6.2	71.2
4	Diabetes mellitus (E10-E14)	393	5.3	61.5
5	Chronic lower respiratory diseases (J40-J47)	235	3.2	36.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	224	3.0	35.1
7	Influenza and pneumonia (J09-J18)	174	2.4	27.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	172	2.3	26.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	109	1.5	17.1
10	Septicemia (A40-A41)	103	1.4	16.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	91	1.2	14.2
12	Parkinson disease (G20-G21)	79	1.1	12.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	1.0	11.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.7	8.4
14	Aortic aneurysm and dissection (I71)	54	0.7	8.4
...	All other causes (residual)	928	12.6	145.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 75-84 years				
...	All causes	9,098	100.0	3,076.9
1	Malignant neoplasms (C00-C97)	2,396	26.3	810.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,102	23.1	710.9
3	Cerebrovascular diseases (I60-I69)	664	7.3	224.6
4	Diabetes mellitus (E10-E14)	469	5.2	158.6
5	Chronic lower respiratory diseases (J40-J47)	410	4.5	138.7
6	Influenza and pneumonia (J09-J18)	320	3.5	108.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	275	3.0	93.0
8	Alzheimer disease (G30)	219	2.4	74.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	218	2.4	73.7
10	Parkinson disease (G20-G21)	206	2.3	69.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	159	1.7	53.8
12	Septicemia (A40-A41)	126	1.4	42.6
13	Pneumonitis due to solids and liquids (J69)	101	1.1	34.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.8	24.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	64	0.7	21.6
...	All other causes (residual)	1,297	14.3	438.6
Non-Hispanic, Asian or Pacific Islander, male, 85 years and over				
...	All causes	9,169	100.0	9,252.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,389	26.1	2,410.8
2	Malignant neoplasms (C00-C97)	1,425	15.5	1,438.0
3	Cerebrovascular diseases (I60-I69)	766	8.4	773.0
4	Alzheimer disease (G30)	529	5.8	533.8
5	Influenza and pneumonia (J09-J18)	507	5.5	511.6
6	Chronic lower respiratory diseases (J40-J47)	457	5.0	461.2
7	Diabetes mellitus (E10-E14)	301	3.3	303.7
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	245	2.7	247.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	236	2.6	238.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	227	2.5	229.1
11	Parkinson disease (G20-G21)	191	2.1	192.7
12	Pneumonitis due to solids and liquids (J69)	154	1.7	155.4
13	Septicemia (A40-A41)	118	1.3	119.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.6	59.5
15	Nutritional deficiencies (E40-E64)	44	0.5	44.4
...	All other causes (residual)	1,521	16.6	1,534.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, all ages				
...	All causes	35,362	100.0	335.6
1	Malignant neoplasms (C00-C97)	9,014	25.5	85.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,072	20.0	67.1
3	Cerebrovascular diseases (I60-I69)	2,988	8.4	28.4
4	Alzheimer disease (G30)	1,793	5.1	17.0
5	Diabetes mellitus (E10-E14)	1,435	4.1	13.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,182	3.3	11.2
7	Influenza and pneumonia (J09-J18)	1,131	3.2	10.7
8	Chronic lower respiratory diseases (J40-J47)	865	2.4	8.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	818	2.3	7.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	745	2.1	7.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	424	1.2	4.0
12	Septicemia (A40-A41)	419	1.2	4.0
13	Parkinson disease (G20-G21)	380	1.1	3.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	257	0.7	2.4
15	Pneumonitis due to solids and liquids (J69)	232	0.7	2.2
...	All other causes (residual)	6,607	18.7	62.7
Non-Hispanic, Asian or Pacific Islander, female, 1-4 years				
...	All causes	52	100.0	11.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	19.2	*
2	Malignant neoplasms (C00-C97)	7	13.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	13.5	*
4	Anemias (D50-D64)	3	5.8	*
4	Influenza and pneumonia (J09-J18)	3	5.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	2	3.8	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	3.8	*
9	Septicemia (A40-A41)	1	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.9	*
9	Hernia (K40-K46)	1	1.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.9	*
...	All other causes (residual)	12	23.1	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 5-14 years				
...	All causes	94	100.0	7.9
1	Malignant neoplasms (C00-C97)	19	20.2	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	20.2	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	11.7	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	9.6	*
5	Septicemia (A40-A41)	4	4.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.1	*
6	Cerebrovascular diseases (I60-I69)	2	2.1	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.1	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.1	*
10	Anemias (D50-D64)	1	1.1	*
10	Diabetes mellitus (E10-E14)	1	1.1	*
10	Acute bronchitis and bronchiolitis (J20-J21)	1	1.1	*
...	All other causes (residual)	21	22.3	1.8
Non-Hispanic, Asian or Pacific Islander, female, 15-24 years				
...	All causes	293	100.0	22.2
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	29.7	6.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	27.3	6.1
3	Malignant neoplasms (C00-C97)	31	10.6	2.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	5.1	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	3.8	*
6	Influenza and pneumonia (J09-J18)	4	1.4	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	4	1.4	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.0	*
9	Diabetes mellitus (E10-E14)	2	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.7	*
12	Septicemia (A40-A41)	1	0.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
12	Meningitis (G00,G03)	1	0.3	*
12	Pneumonitis due to solids and liquids (J69)	1	0.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (residual)	44	15.0	3.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 25-34 years				
...	All causes	487	100.0	27.4
1	Malignant neoplasms (C00-C97)	126	25.9	7.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	17.2	4.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	15.2	4.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	5.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.9	1.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	18	3.7	*
7	Cerebrovascular diseases (I60-I69)	14	2.9	*
8	Influenza and pneumonia (J09-J18)	7	1.4	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.4	*
10	Diabetes mellitus (E10-E14)	5	1.0	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.8	*
14	Septicemia (A40-A41)	3	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
...	All other causes (residual)	84	17.2	4.7
Non-Hispanic, Asian or Pacific Islander, female, 35-44 years				
...	All causes	853	100.0	50.6
1	Malignant neoplasms (C00-C97)	352	41.3	20.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	8.3	4.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	8.0	4.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	7.5	3.8
5	Cerebrovascular diseases (I60-I69)	53	6.2	3.1
6	Diabetes mellitus (E10-E14)	25	2.9	1.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	19	2.2	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	2.2	*
9	Septicemia (A40-A41)	15	1.8	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	13	1.5	*
11	Influenza and pneumonia (J09-J18)	10	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.2	*
13	Chronic lower respiratory diseases (J40-J47)	9	1.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.6	*
...	All other causes (residual)	115	13.5	6.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 45-54 years				
...	All causes	1,891	100.0	130.0
1	Malignant neoplasms (C00-C97)	989	52.3	68.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	180	9.5	12.4
3	Cerebrovascular diseases (I60-I69)	121	6.4	8.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	5.3	6.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	3.6	4.7
6	Diabetes mellitus (E10-E14)	68	3.6	4.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	24	1.3	1.7
8	Septicemia (A40-A41)	18	1.0	*
9	Influenza and pneumonia (J09-J18)	17	0.9	*
10	Chronic lower respiratory diseases (J40-J47)	16	0.8	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.8	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	0.8	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	0.5	*
...	All other causes (residual)	221	11.7	15.2
Non-Hispanic, Asian or Pacific Islander, female, 55-64 years				
...	All causes	3,471	100.0	291.9
1	Malignant neoplasms (C00-C97)	1,740	50.1	146.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	463	13.3	38.9
3	Cerebrovascular diseases (I60-I69)	202	5.8	17.0
4	Diabetes mellitus (E10-E14)	155	4.5	13.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	3.3	9.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	1.8	5.2
7	Influenza and pneumonia (J09-J18)	56	1.6	4.7
8	Chronic lower respiratory diseases (J40-J47)	51	1.5	4.3
9	Septicemia (A40-A41)	46	1.3	3.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	42	1.2	3.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	42	1.2	3.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	1.2	3.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.7	1.9
14	Viral hepatitis (B15-B19)	22	0.6	1.8
15	Aortic aneurysm and dissection (I71)	19	0.5	*
...	All other causes (residual)	393	11.3	33.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 65-74 years				
...	All causes	5,541	100.0	691.2
1	Malignant neoplasms (C00-C97)	2,133	38.5	266.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	979	17.7	122.1
3	Cerebrovascular diseases (I60-I69)	419	7.6	52.3
4	Diabetes mellitus (E10-E14)	281	5.1	35.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	158	2.9	19.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	120	2.2	15.0
7	Influenza and pneumonia (J09-J18)	116	2.1	14.5
8	Chronic lower respiratory diseases (J40-J47)	106	1.9	13.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	87	1.6	10.9
10	Septicemia (A40-A41)	75	1.4	9.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	70	1.3	8.7
12	Parkinson disease (G20-G21)	61	1.1	7.6
13	Alzheimer disease (G30)	58	1.0	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	0.7	5.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.6	4.2
...	All other causes (residual)	803	14.5	100.2
Non-Hispanic, Asian or Pacific Islander, female, 75-84 years				
...	All causes	8,432	100.0	2,203.0
1	Malignant neoplasms (C00-C97)	2,058	24.4	537.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,788	21.2	467.1
3	Cerebrovascular diseases (I60-I69)	786	9.3	205.4
4	Diabetes mellitus (E10-E14)	417	4.9	108.9
5	Alzheimer disease (G30)	339	4.0	88.6
6	Chronic lower respiratory diseases (J40-J47)	278	3.3	72.6
7	Influenza and pneumonia (J09-J18)	259	3.1	67.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	246	2.9	64.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	227	2.7	59.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	214	2.5	55.9
11	Parkinson disease (G20-G21)	142	1.7	37.1
12	Septicemia (A40-A41)	118	1.4	30.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	56	0.7	14.6
14	Pneumonitis due to solids and liquids (J69)	52	0.6	13.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	51	0.6	13.3
...	All other causes (residual)	1,401	16.6	366.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 85 years and over				
...	All causes	13,819	100.0	8,166.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,539	25.6	2,091.5
2	Malignant neoplasms (C00-C97)	1,555	11.3	919.0
3	Cerebrovascular diseases (I60-I69)	1,388	10.0	820.3
4	Alzheimer disease (G30)	1,387	10.0	819.7
5	Influenza and pneumonia (J09-J18)	657	4.8	388.3
6	Diabetes mellitus (E10-E14)	481	3.5	284.3
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	457	3.3	270.1
8	Chronic lower respiratory diseases (J40-J47)	398	2.9	235.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	309	2.2	182.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	283	2.0	167.2
11	Parkinson disease (G20-G21)	165	1.2	97.5
12	Pneumonitis due to solids and liquids (J69)	138	1.0	81.6
13	Septicemia (A40-A41)	134	1.0	79.2
14	Nutritional deficiencies (E40-E64)	69	0.5	40.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.5	40.2
...	All other causes (residual)	2,791	20.2	1,649.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017